



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/82, 5/04, 15/52, 9/00, C07K 14/415, A01H 1/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 00/42205</b>
			(43) International Publication Date: <b>20 July 2000 (20.07.00)</b>
(21) International Application Number: <b>PCT/EP00/00246</b>		27613 (US). WEGRICH, Lynette, Marcia [US/US]; 112 Windbyrne Drive, Cary, NC 27513 (US).	
(22) International Filing Date: <b>13 January 2000 (13.01.00)</b>		(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).	
(30) Priority Data:		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
09/232,760 15 January 1999 (15.01.99) US 09/237,479 26 January 1999 (26.01.99) US 09/244,288 3 February 1999 (03.02.99) US 09/252,336 18 February 1999 (18.02.99) US 09/281,376 30 March 1999 (30.03.99) US			
(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).			
(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT MBH . [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): LEVIN, Joshua, Zvi [US/US]; 1008 Urban Avenue, Durham, NC 27701 (US). BUDZISZEWSKI, Gregory, Joseph [US/US]; 2016 Englewood Avenue, Durham, NC 27705 (US). POTTER, Sharon, Lee [US/US]; 3837 Whispering Branch Road, Raleigh, NC		<b>Published</b> Without international search report and to be republished upon receipt of that report.	
(54) Title: <b>HERBICIDE TARGET GENE AND METHODS</b>			
(57) Abstract			
<p>The invention relates to genes isolated from <i>Arabidopsis</i> that code for proteins essential for seedling growth. The invention also includes the methods of using these proteins to discover new herbicides, based on the essentiality of the genes for normal growth and development. The invention can also be used in a screening assay to identify inhibitors that are potential herbicides. The invention is also applied to the development of herbicide tolerant plants, plant tissues, plant seeds, and plant cells.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## HERBICIDE TARGET GENE AND METHODS

The invention relates to genes isolated from *Arabidopsis* that code for proteins essential for seedling growth. The invention also includes the methods of using these proteins as an herbicide target, based on the essentiality of the gene for normal growth and development. The invention is also useful as a screening assay to identify inhibitors that are potential herbicides. The invention may also be applied to the development of herbicide tolerant plants, plant tissues, plant seeds, and plant cells.

The use of herbicides to control undesirable vegetation such as weeds in crop fields has become almost a universal practice. The herbicide market exceeds 15 billion dollars annually. Despite this extensive use, weed control remains a significant and costly problem for farmers.

Effective use of herbicides requires sound management. For instance, the time and method of application and stage of weed plant development are critical to getting good weed control with herbicides. Since various weed species are resistant to herbicides, the production of effective new herbicides becomes increasingly important. Novel herbicides can now be discovered using high-throughput screens that implement recombinant DNA technology. Metabolic enzymes found to be essential to plant growth and development can be recombinantly produced through standard molecular biological techniques and utilized as herbicide targets in screens for novel inhibitors of the enzyme activity. The novel inhibitors discovered through such screens may then be used as herbicides to control undesirable vegetation.

Herbicides that exhibit greater potency, broader weed spectrum, and more rapid degradation in soil can also, unfortunately, have greater crop phytotoxicity. One solution applied to this problem has been to develop crops that are resistant or tolerant to herbicides. Crop hybrids or varieties tolerant to the herbicides allow for the use of the herbicides to kill weeds without attendant risk of damage to the crop. Development of tolerance can allow application of a herbicide to a crop where its use was previously precluded or limited (e.g. to pre-emergence use) due to sensitivity of the crop to the herbicide. For example, U.S. Patent No. 4,761,373 to Anderson *et al.* is directed to plants resistant to various imidazollnone or sulfonamide herbicides. An altered acetohydroxyacid synthase (AHAS) enzyme confers the resistance. U.S. Patent No. 4,975,374 to Goodman *et al.* relates to plant cells and plants containing a gene encoding a mutant glutamine

synthetase (GS) resistant to inhibition by herbicides that were known to inhibit GS, *e.g.* phosphinothricin and methionine sulfoximine. U.S. Patent No. 5,013,659 to Bedbrook *et al.* is directed to plants expressing a mutant acetolactate synthase that renders the plants resistant to inhibition by sulfonylurea herbicides. U.S. Patent No. 5,162,602 to Somers *et al.* discloses plants tolerant to inhibition by cyclohexanedione and aryloxyphenoxypropanoic acid herbicides. The tolerance is conferred by an altered acetyl coenzyme A carboxylase (ACCase).

Notwithstanding the above described advancements, there remain persistent and ongoing problems with unwanted or detrimental vegetation growth (*e.g.* weeds). Furthermore, as the population continues to grow, there will be increasing food shortages. Therefore, there exists a long felt, yet unfulfilled need, to find new, effective, and economic herbicides.

For clarity, certain terms used in the specification are defined and presented as follows:

Chimeric: is used to indicate that a DNA sequence, such as a vector or a gene, is comprised of more than one DNA sequences of distinct origin which are fused together by recombinant DNA techniques resulting in a DNA sequence, which does not occur naturally, and which particularly does not occur in the plant to be transformed.

Co-factor: natural reactant, such as an organic molecule or a metal ion, required in an enzyme-catalyzed reaction. A co-factor is *e.g.* NAD(P), riboflavin (including FAD and FMN), folate, molybdopterin, thiamin, biotin, lipoic acid, pantothenic acid and coenzyme A, S-adenosylmethionine, pyridoxal phosphate, ubiquinone, menaquinone. Optionally, a co-factor can be regenerated and reused.

DNA shuffling: DNA shuffling is a method to rapidly, easily and efficiently introduce mutations or rearrangements, preferably randomly, in a DNA molecule or to generate exchanges of DNA sequences between two or more DNA molecules, preferably randomly. The DNA molecule resulting from DNA shuffling is a shuffled DNA molecule that is a non-naturally occurring DNA molecule derived from at least one template DNA molecule. The shuffled DNA encodes an enzyme modified with respect to the enzyme encoded by the template DNA, and preferably has an altered biological activity with respect to the enzyme encoded by the template DNA.

Enzyme activity: means herein the ability of an enzyme to catalyze the conversion of a substrate into a product. A substrate for the enzyme comprises the natural substrate of the enzyme but also comprises analogues of the natural substrate, which can also be



converted, by the enzyme into a product or into an analogue of a product. The activity of the enzyme is measured for example by determining the amount of product in the reaction after a certain period of time, or by determining the amount of substrate remaining in the reaction mixture after a certain period of time. The activity of the enzyme is also measured by determining the amount of an unused co-factor of the reaction remaining in the reaction mixture after a certain period of time or by determining the amount of used co-factor in the reaction mixture after a certain period of time. The activity of the enzyme is also measured by determining the amount of a donor of free energy or energy-rich molecule (e.g. ATP, phosphoenolpyruvate, acetyl phosphate or phosphocreatine) remaining in the reaction mixture after a certain period of time or by determining the amount of a used donor of free energy or energy-rich molecule (e.g. ADP, pyruvate, acetate or creatine) in the reaction mixture after a certain period of time.

Expression: refers to the transcription and/or translation of an endogenous gene or a transgene in plants. In the case of antisense constructs, for example, expression may refer to the transcription of the antisense DNA only.

Gene: refers to a coding sequence and associated regulatory sequences wherein the coding sequence is transcribed into RNA such as mRNA, rRNA, tRNA, snRNA, sense RNA or antisense RNA. Examples of regulatory sequences are promoter sequences, 5' and 3' untranslated sequences and termination sequences. Further elements that may be present are, for example, introns.

Herbicide: a chemical substance used to kill or suppress the growth of plants, plant cells, plant seeds, or plant tissues.

Heterologous DNA Sequence: a DNA sequence not naturally associated with a host cell into which it is introduced, including non-naturally occurring multiple copies of a naturally occurring DNA sequence; and genetic constructs wherein an otherwise homologous DNA sequence is operatively linked to a non-native sequence.

Homologous DNA Sequence: a DNA sequence naturally associated with a host cell into which it is introduced.

Inhibitor: a chemical substance that causes abnormal growth, e.g., by inactivating the enzymatic activity of a protein such as a biosynthetic enzyme, receptor, signal transduction protein, structural gene product, or transport protein that is essential to the growth or survival of the plant. In the context of the instant invention, an inhibitor is a chemical substance that alters the enzymatic activity encoded by the 245 gene, the 5283 gene, the 2490 gene, the 3963 gene or the 4036 gene from a plant. More generally, an

inhibitor causes abnormal growth of a host cell by interacting with the gene product encoded by the 245 gene, the 5283 gene, the 2490 gene, the 3963 gene or the 4036 gene.

Isogenic: plants which are genetically identical, except that they may differ by the presence or absence of a heterologous DNA sequence.

Isolated: in the context of the present invention, an isolated DNA molecule or an isolated enzyme is a DNA molecule or enzyme that, by the hand of man, exists apart from its native environment and is therefore not a product of nature. An isolated DNA molecule or enzyme may exist in a purified form or may exist in a non-native environment such as, for example, in a transgenic host cell.

Marker gene: a gene encoding a selectable or screenable trait

Mature protein: protein which is normally targeted to a cellular organelle, such as a chloroplast, and from which the transit peptide has been removed.

Minimal Promoter: promoter elements, particularly a TATA element, that are inactive or that have greatly reduced promoter activity in the absence of upstream activation. In the presence of a suitable transcription factor, the minimal promoter functions to permit transcription.

Modified Enzyme Activity: enzyme activity different from that which naturally occurs in a plant (i.e. enzyme activity that occurs naturally in the absence of direct or indirect manipulation of such activity by man), which is tolerant to inhibitors that inhibit the naturally occurring enzyme activity.

Plant: refers to any plant, particularly to seed plants

Plant cell: structural and physiological unit of the plant, comprising a protoplast and a cell wall. The plant cell may be in form of an isolated single cell or a cultured cell, or as a part of higher organized unit such as, for example, a plant tissue, or a plant organ.

Plant material: refers to leaves, stems, roots, flowers or flower parts, fruits, pollen, pollen tubes, ovules, embryo sacs, egg cells, zygotes, embryos, seeds, cuttings, cell or tissue cultures, or any other part or product of a plant

Pre-protein: protein which is normally targeted to a cellular organelle, such as a chloroplast, and still comprising its transit peptide.

Recombinant DNA molecule: a combination of DNA sequences that are joined together using recombinant DNA technology

Selectable marker gene: a gene whose expression does not confer a selective advantage to a transformed cell, but whose expression makes the transformed cell phenotypically distinct from untransformed cells.

Significant Increase: an increase in enzymatic activity that is larger than the margin of error inherent in the measurement technique, preferably an increase by about 2-fold or greater of the activity of the wild-type enzyme in the presence of the inhibitor, more preferably an increase by about 5-fold or greater, and most preferably an increase by about 10-fold or greater.

Significantly less: means that the amount of a product of an enzymatic reaction is reduced by more than the margin of error inherent in the measurement technique, preferably a decrease by about 2-fold or greater of the activity of the wild-type enzyme in the absence of the inhibitor, more preferably an decrease by about 5-fold or greater, and most preferably an decrease by about 10-fold or greater.

In its broadest sense, the term "substantially similar", when used herein with respect to a nucleotide sequence, means a nucleotide sequence corresponding to a reference nucleotide sequence, wherein the corresponding sequence encodes a polypeptide having substantially the same structure and function as the polypeptide encoded by the reference nucleotide sequence, e.g. where only changes in amino acids not affecting the polypeptide function occur. Desirably the substantially similar nucleotide sequence encodes the polypeptide encoded by the reference nucleotide sequence. The term "substantially similar" is specifically intended to include nucleotide sequences wherein the sequence has been modified to optimize expression in particular cells. The percentage of identity between the substantially similar nucleotide sequence and the reference nucleotide sequence desirably is at least 65%, more desirably at least 75%, preferably at least 85%, more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. Sequence comparisons are carried out using a Smith-Waterman sequence alignment algorithm (see e.g. Waterman, M.S. Introduction to Computational Biology: Maps, sequences and genomes. Chapman & Hall. London: 1995. ISBN 0-412-99391-0.). The localS program, version 1.16, is used with following parameters: match: 1, mismatch penalty: 0.33, open-gap penalty: 2, extended-gap penalty: 2. A nucleotide sequence "substantially similar" to reference nucleotide sequence hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at

50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO<sub>4</sub>, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C. As used herein the term "245 gene", "5283 gene", "2490 gene", "3963 gene" or "4036 gene" refers to a DNA molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, respectively, or comprising a nucleotide sequence substantially similar to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9, respectively. Homologs of the 245 gene, the 5283 gene, the 2490 gene, the 3963 gene or the 4036 gene include nucleotide sequences that encode an amino acid sequence that is at least 30% identical to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10, respectively, as measured, using the parameters described below, wherein the amino acid sequence encoded by the homolog has the biological activity of the 245, 5283, 2490, 3963, or 4036 protein, respectively.

The term "substantially similar", when used herein with respect to a protein, means a protein corresponding to a reference protein, wherein the protein has substantially the same structure and function as the reference protein, e.g. where only changes in amino acids sequence not affecting the polypeptide function occur. When used for a protein or an amino acid sequence the percentage of identity between the substantially similar and the reference protein or amino acid sequence desirably is at least 65%, more desirably at least 75%, preferably at least 85%, more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%, using default BLAST analysis parameters. As used herein the term "245 protein", "5283 protein", "2490 protein", "3963 protein" or "4036 protein" refers to an amino acid sequence encoded by a DNA molecule comprising a nucleotide sequence substantially similar to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, respectively. Homologs of the 245 protein, the 5283 protein, the 2490 protein, the 3963 protein or the 4036 protein are amino acid sequences that are at least 30% identical to SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10, respectively, as measured using the parameters described below, wherein the homologs have the biological activity of the 245, 5283, 2490, 3963, or 4036 protein, respectively.

One skilled in the art is also familiar with other analysis tools, such as GAP analysis, to determine the percentage of identity between the "substantially similar" and the reference nucleotide sequence, or protein or amino acid sequence. In the present invention, "substantially similar" is therefore also determined using default GAP analysis parameters with the University of Wisconsin GCG, SEQWEB application of GAP, based on the algorithm of Needleman and Wunsch (Needleman and Wunsch (1970) J Mol. Biol. 48: 443-

453).

Thus, in the context of the "245 gene" and using GAP analysis as described above, "substantially similar" refers to nucleotide sequences that encode a protein having at least 47% identity, more preferably at least 60% identity, still more preferably at least 75% identity, still more preferably at least 85% identity, still more preferably at least 95% identity, yet still more preferably at least 99% identity to SEQ ID NO:2.

In the context of the "5283 gene" and using GAP analysis as described above, "substantially similar" refers to nucleotide sequences that encode a protein having at least 74% identity, more preferably at least 80% identity, still more preferably at least 85% identity, still more preferably at least 90% identity, still more preferably at least 95% identity, yet still more preferably at least 99% identity to SEQ ID NO:4. Also, "substantially similar" preferably also refers to nucleotide sequences having at least 80% identity, more preferably at least 90% identity, still more preferably 95% identity, yet still more preferably at least 99% identity, to SEQ ID NO:3, wherein said nucleotide sequence comparisons are conducted using GAP analysis as described above.

In the context of the "2490 gene" and using GAP analysis as described above, "substantially similar" refers to nucleotide sequences that encode a protein having at least 82% identity, more preferably at least 85% identity, more preferably at least 90% identity, still more preferably at least 95% identity, yet still more preferably at least 99% identity to SEQ ID NO:6. Also, "substantially similar" preferably also refers to nucleotide sequences having at least 87% identity, more preferably at least 90% identity, still more preferably 95% identity, yet still more preferably at least 99% identity, to SEQ ID NO:5, wherein said nucleotide sequence comparisons are conducted using GAP analysis as described above.

In the context of the "3963 gene" and using GAP analysis as described above, "substantially similar" refers to nucleotide sequences that encode a protein having at least 40% identity, more preferably at least 60% identity, more preferably at least 80% identity, still more preferably at least 90% identity, still more preferably at least 95% identity, yet still more preferably at least 99% identity to SEQ ID NO:8. Also, "substantially similar" preferably also refers to nucleotide sequences having at least 49% identity, more preferably at least 60% identity, still more preferably 80% identity, more preferably at least 90% identity, more preferably at least 95% identity, yet still more preferably at least 99% identity, to SEQ ID NO:7, wherein said nucleotide sequence comparisons are conducted using GAP analysis as described above.

In the context of the "4036 gene" and using GAP analysis as described above, "substantially similar" refers to nucleotide sequences that encode a protein having at least 67% identity, more preferably at least 80% identity, more preferably at least 85% identity, still more preferably at least 90% identity, still more preferably at least 95% identity, yet still more preferably at least 99% identity to SEQ ID NO:10.

Further, using GAP analysis as described above, "homologs of the 245 gene" include nucleotide sequences that encode an amino acid sequence that has at least 24% identity to SEQ ID NO:2, more preferably at least 30% identity, still more preferably at least 40% identity, still more preferably at least 45% identity, yet still more preferably at least 55% identity, still more preferably at least 65% identity, yet still more preferably at least 75% identity to SEQ ID NO:2, wherein the amino acid sequence encoded by the homolog has the biological activity of the 245 protein.

Further, using GAP analysis as described above, "homologs of the 5283 gene" include nucleotide sequences that encode an amino acid sequence that has at least 23% identity to SEQ ID NO:4, more preferably at least 40% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 74% identity to SEQ ID NO:4, wherein the amino acid sequence encoded by the homolog has the biological activity of the 5283 protein.

Further, using GAP analysis as described above, "homologs of the 2490 gene" include nucleotide sequences that encode an amino acid sequence that has at least 30% identity to SEQ ID NO:6, more preferably at least 30% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 80% identity to SEQ ID NO:6, wherein the amino acid sequence encoded by the homolog has the biological activity of the 2490 protein.

Further, using GAP analysis as described above, "homologs of the 3963 gene" include nucleotide sequences that encode an amino acid sequence that has at least 34% identity to SEQ ID NO:8, more preferably at least 40% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 75% identity to SEQ ID NO:8, wherein the amino acid sequence encoded by the homolog has the biological activity of the 3963 protein.

Further, using GAP analysis as described above, "homologs of the 4036 gene" include nucleotide sequences that encode an amino acid sequence that has at least 44% identity to SEQ ID NO:10, more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 75% identity to SEQ ID NO:10, wherein the

amino acid sequence encoded by the homolog has the biological activity of the 4036 protein.

When using GAP analysis as described above with respect to a protein or an amino acid sequence and in the context of the "245 gene", the percentage of identity between the "substantially similar" protein or amino acid sequence and the reference protein or amino acid sequence (in this case SEQ ID NO:2) is at least 47%, more preferably at least 60%, still more preferably at least 75%, still more preferably at least 85%, still more preferably at least 95%, yet still more preferably at least 99%. "Homologs of the 245 protein" include amino acid sequences that are at least 24% identical to SEQ ID NO:2, more preferably at least 30% identical, still more preferably at least 40% identical, still more preferably at least 45% identical, yet still more preferably at least 55% identical, still more preferably at least 65% identical, yet still more preferably at least 75% identical to SEQ ID NO:2, wherein homologs of the 245 protein have the biological activity of the 245 protein.

In the context of the "5283 gene" and using GAP analysis as described above, the percentage of identity between the substantially similar protein or amino acid sequence and the reference protein or amino acid sequence (in this case SEQ ID NO:4) is at least 74%, more preferably at least 80%, still more preferably at least 85%, still more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. "Homologs of the 5283 protein" include amino acid sequences that at least 23% identity to SEQ ID NO:4, more preferably at least 40% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 74% identity to SEQ ID NO:4, wherein homologs of the 5283 protein have the biological activity of the 5283 protein.

In the context of the "2490 gene" and using GAP analysis as described above, the percentage of identity between the substantially similar protein or amino acid sequence and the reference protein or amino acid sequence (in this case SEQ ID NO:6) is at least 82%, more preferably at least 85%, more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. "Homologs of the 2490 protein" include amino acid sequences that have at least 30% identity to SEQ ID NO:6, more preferably at least 30% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 80% identity to SEQ ID NO:6, wherein the homologs of the 2490 protein have the biological activity of the 2490 protein.

In the context of the "3963 gene" and using GAP analysis as described above, the percentage of identity between the substantially similar protein or amino acid sequence and the reference protein or amino acid sequence (in this case SEQ ID NO:8) is at least 40%,

more preferably at least 60%, more preferably at least 80%, still more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. "Homologs of the 3963 protein" include amino acid sequences that has at least 34% identity to SEQ ID NO:8, more preferably at least 40% identity, still more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 75% identity to SEQ ID NO:8, wherein the homologs of the 3963 protein have the biological activity of the 3963 protein.

In the context of the "4036 gene" and using GAP analysis as described above, the percentage of identity between the substantially similar reference protein or amino acid sequence and the reference protein or amino acid sequence (in this case SEQ ID NO:10) is at least 67%, more preferably at least 80%, more preferably at least 85%, still more preferably at least 90%, still more preferably at least 95%, yet still more preferably at least 99%. "Homologs of the 4036 protein" include amino acid sequences that have at least 44% identity to SEQ ID NO:10, more preferably at least 50% identity, still more preferably at least 60% identity, yet still more preferably at least 75% identity to SEQ ID NO:10, wherein the homologs of the 4036 protein has the biological activity of the 4036 protein.

Substrate: a substrate is the molecule that an enzyme naturally recognizes and converts to a product in the biochemical pathway in which the enzyme naturally carries out its function, or is a modified version of the molecule, which is also recognized by the enzyme and is converted by the enzyme to a product in an enzymatic reaction similar to the naturally-occurring reaction.

Tolerance: the ability to continue essentially normal growth or function when exposed to an inhibitor or herbicide in an amount sufficient to suppress the normal growth or function of native, unmodified plants.

Transformation: a process for introducing heterologous DNA into a cell, tissue, or plant. Transformed cells, tissues, or plants are understood to encompass not only the end product of a transformation process, but also transgenic progeny thereof.

Transgenic: stably transformed with a recombinant DNA molecule that preferably comprises a suitable promoter operatively linked to a DNA sequence of interest.

SEQ ID NO:1 cDNA sequence for the Arabidopsis 245 gene

SEQ ID NO:2 amino acid sequence encoded by the Arabidopsis 245 DNA sequence shown in SEQ ID NO:1



SEQ ID NO:3 cDNA sequence for the Arabidopsis 5283 gene  
SEQ ID NO:4 amino acid sequence encoded by the Arabidopsis 5283 DNA sequence shown in SEQ ID NO:3  
SEQ ID NO:5 cDNA sequence for the Arabidopsis 2490 gene  
SEQ ID NO:6 amino acid sequence encoded by the Arabidopsis 2490 DNA sequence shown in SEQ ID NO:5  
SEQ ID NO:7 cDNA sequence for the Arabidopsis 3963 gene  
SEQ ID NO:8 amino acid sequence encoded by the Arabidopsis 3963 DNA sequence shown in SEQ ID NO:7  
SEQ ID NO:9 cDNA sequence for the Arabidopsis 4036 gene  
SEQ ID NO:10 amino acid sequence encoded by the Arabidopsis 4036 DNA sequence shown in SEQ ID NO:9  
SEQ ID NO:11 oligonucleotide SLP346for  
SEQ ID NO:12 partial genomic sequence of the Arabidopsis 245 gene  
SEQ ID NO:13 3'UTR from the cDNA sequence for the Arabidopsis 245 gene  
SEQ ID NO:14 genomic sequence of the Arabidopsis 5283 gene  
SEQ ID NO:15 oligonucleotide SLP328  
SEQ ID NO:16 oligonucleotide LW60  
SEQ ID NO:17 5'UTR from the cDNA sequence for the Arabidopsis 5283 gene  
SEQ ID NO:18 3'UTR from the cDNA sequence for the Arabidopsis 5283 gene  
SEQ ID NO:19 genomic sequence of the Arabidopsis 2490 gene  
SEQ ID NO:20 5'UTR from the cDNA for the Arabidopsis 2490 gene  
SEQ ID NO:21 3'UTR from the cDNA sequence for the Arabidopsis 2490 gene  
SEQ ID NO:22 oligonucleotide SLP369  
SEQ ID NO:23 oligonucleotide SLP370  
SEQ ID NO:24 genomic sequence of the Arabidopsis 3963 gene  
SEQ ID NO:25 oligonucleotide -21  
SEQ ID NO:26 3'UTR from the cDNA sequence for the Arabidopsis 3963 gene  
SEQ ID NO:27 genomic sequence of the Arabidopsis 4036 gene  
SEQ ID NO:28 cDNA coding sequence for the Arabidopsis 4036 gene including variations between the cDNA and genomic sequence from cultivars Landsberg and Columbia  
SEQ ID NO:29 amino acid sequence encoded by the Arabidopsis 4036 DNA sequence shown in SEQ ID NO:28

Encompassed by the invention is an isolated DNA molecule comprising a nucleotide sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. Preferred is the DNA molecule according to the invention, wherein the sequence encodes an amino acid sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. Further preferred is DNA molecule according to the invention, wherein the sequence is any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. Further preferred is the DNA molecule according to the invention, wherein the sequence encodes the amino acid sequence of any one of the sequences selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. Further preferred is a DNA molecule according to the invention, wherein said nucleotide sequence is a plant nucleotide sequence. More preferred is the DNA molecule according to the invention, wherein the plant is *Arabidopsis thaliana*. Further preferred is a DNA molecule according to the invention, wherein the protein has any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 396 activity and 4036 activity. Further encompassed by the invention is an amino acid sequence comprising an amino acid sequence encoded by a nucleotide sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. Preferred is the amino acid sequence according to the invention comprising an amino acid sequence encoded by any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. A further object of the invention is an amino acid sequence comprising an amino acid sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. Preferred is the amino acid sequence according to the invention, wherein the sequence is any one of the sequences selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. Further preferred is the amino acid sequence according to the invention, wherein the protein has any one of the activities selected from the group consisting of 245, 5283, 2490, 3963 and 4036 activity. Encompassed by the invention is an amino acid sequence comprising at least 20 consecutive amino acid residues of the amino acid sequence encoded by any one of the

sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. Further encompassed is an amino acid sequence comprising at least 20 consecutive amino acid residues of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10. An object of the invention is an expression cassette comprising a promoter operatively linked to a DNA molecule according to the invention. Further encompassed by the invention is a recombinant vector comprising an expression cassette according to the invention, wherein said vector is capable of being stably transformed into a host cell. Further encompassed is a host cell comprising an expression cassette according to the invention, wherein said nucleotide sequence is expressible in said cell. Preferred is a host cell according to the invention, wherein said host cell is an eukaryotic cell. More preferred is a host cell according to the invention, wherein said host cell is selected from the group consisting of an insect cell, a yeast cell, and a plant cell. Also more preferred is a host cell according to the invention, wherein said host cell is a prokaryotic cell. Also more preferred is a host cell according to the invention, wherein said host cell is a bacterial cell. Encompassed is a plant or seed comprising a plant cell according to the invention. Preferred is a plant according to the invention, wherein said plant is tolerant to an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity.

Further encompassed in the invention is a method comprising obtaining a host cell comprising a heterologous DNA molecule encoding a protein having 245, 5283, 2490, 3963, or 4036 activity; and expressing said protein in said host cell. Preferably the host cell is a bacterial cell, a yeast cell or an insect cell.

Further encompassed is a process for making nucleotides sequences encoding gene products having altered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity comprising,

- a) shuffling a nucleotide sequence of claim 1,
- b) expressing the resulting shuffled nucleotide sequences and
- c) selecting for altered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity as compared to the activity selected

from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity of the gene product of said unmodified nucleotide sequence.

Preferred is a process according to the invention, wherein the nucleotide sequence is any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. Encompassed by the invention is a shuffled DNA molecule obtainable by the process according to the invention. Encompassed by the invention is a shuffled DNA molecule produced by the process according to the invention. Further encompassed by the invention is a shuffled DNA molecule obtained by the process according to the invention, wherein said shuffled DNA molecule encodes a gene product having enhanced tolerance to an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity. A further object of the invention is an expression cassette comprising a promoter operatively linked to a nucleotide sequence according to the invention. Further encompassed by the invention is a recombinant vector comprising an expression cassette according to the invention, wherein said vector is capable of being stably transformed into a host cell. A further object of the invention is a host cell comprising an expression cassette according to the invention, wherein said nucleotide sequence is expressible in said cell. Preferred is a host cell according to the invention, wherein said host cell is an eukaryotic cell. Also preferred is a host cell according to the invention, wherein said host cell is selected from the group consisting of an insect cell, a yeast cell, and a plant cell. Also preferred is a host cell according to the invention, wherein said host cell is a prokaryotic cell. Also preferred is a host cell according to the invention, wherein said host cell is a bacterial cell. An object of the invention is a plant or seed comprising a plant cell according to the invention. Preferred is a plant according to the invention, wherein said plant is tolerant to an inhibitor selected from the group consisting of 245, 5283, 2490, 3963 and 4036 activity. Further encompassed is a method for selecting compounds that interact with the protein encoded by any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9, comprising:

a) expressing a DNA molecule comprising any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9, respectively, or a sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9 to generate the corresponding protein,

b) testing a compound suspected of having the ability to interact with the protein expressed in step (a), and

c) selecting compounds that interact with the protein in step (b).

A further object of the invention is a process of identifying an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity comprising:

- a) introducing a DNA molecule comprising a nucleotide sequence of any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9, respectively, and having any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity, or nucleotide sequences substantially similar thereto, or a homolog thereof, into a plant cell, such that said sequence is functionally expressible at levels that are higher than wild-type expression levels,
- b) combining said plant cell with a compound to be tested for the ability to inhibit any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity under conditions conducive to such inhibition,
- c) measuring plant cell growth under the conditions of step (b), and
- d) comparing the growth of said plant cell with the growth of a plant cell having an unaltered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity under identical conditions, and
- e) selecting said compound that inhibits plant cell growth in step (d).

Encompassed by the invention is a compound having herbicidal activity identifiable according to the process according to the invention. Further encompassed is a process of identifying compounds having herbicidal activity comprising:

- a) combining a protein according to the invention and a compound to be tested for the ability to interact with said protein, under conditions conducive to interaction,
- b) selecting a compound identified in step (a) that is capable of interacting with said protein,
- c) applying identified compound in step (b) to a plant to test for herbicidal activity, and

- d) selecting compounds having herbicidal activity.

Further encompassed is a compound having herbicidal activity identifiable according to the process according to the invention. A further object of the invention is a method for suppressing the growth of a plant comprising, applying to said plant a compound that inhibits the activity of the amino acid sequence according to the invention in an amount sufficient to suppress the growth of said plant.

Preferred is the method according to the invention, wherein the compound is a compound having herbicidal activity identifiable according to the process according to the invention. Encompassed is a method of improving crops comprising, applying to a herbicide tolerant plant or seed according to the invention, a compound having herbicidal activity identifiable according to a process according to the invention, in an amount that inhibits the growth of undesired vegetation without significantly suppressing the growth of the herbicide tolerant plant or seed. An object of the invention is a DNA molecule comprising a nucleotide sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29.

It is an object of the invention to provide an effective and beneficial method to identify novel herbicides. A feature of the invention is the identification of a gene in *Arabidopsis*, herein referred to as the 245 gene, which shows sequence similarity to peptide release factor 2 (Craigie et al. (1985) Proc. Natl. Acad. Sci. 82: 3616-3620; Craigie and Caskey (1987) Biochimie 69: 1031-1041; Ito et al. (1998) Proc. Natl. Acad. Sci., 95: 8165-8169). Another feature of the invention is the discovery that the 245 gene is essential for seedling growth and development. An advantage of the present invention is that the newly discovered essential gene containing a novel herbicidal mode of action enables one skilled in the art to easily and rapidly identify novel herbicides.

A further feature of the invention is the identification of a gene in *Arabidopsis*, herein referred to as the 5283 gene, which shows sequence similarity to the following: an uncharacterized gene from *Schizosaccharomyces pombe*; the *Saccharomyces cerevisiae* PRP31 gene that encodes a factor essential for pre-mRNA splicing (Weidenhammer et al. (1996) Nucleic Acids Res. 24: 1164-1170; Weidenhammer et al. (1997) Mol. Cell. Biol., 17:

3580-3585); the *Pisum sativum* SARBP-1 and SARBP-2 genes that encode Scaffold Attachment Region (SAR) DNA-binding proteins (Rzepecki et al. (1995) Acta Biochim. Pol., 42: 75-81); and the *Saccharomyces cerevisiae* SIK1 gene that encodes a protein that can suppress the growth inhibitory effects of IKB (Morin et al. (1995) Cell Growth & Differentiation, 6: 789-798). The SIK1 gene product is also referred to as Nop56, which is shown to be an essential nucleolar protein (Gautier et al. (1997) Mol. Cell. Biol. 17: 7088-7098). Another feature of the invention is the discovery that the 5283 gene is essential for seedling growth and development. An advantage of the present invention is that the newly discovered essential gene containing a novel herbicidal mode of action enables one skilled in the art to easily and rapidly identify novel herbicides.

A further feature of the invention is the identification of a gene in *Arabidopsis*, herein referred to as the 2490 gene, which encodes a protein with sequence similarity to a chloroplast envelope protein (Ko et al. (1995) The Journal of Biological Chem. 270: 28601-28608; Wu et al. (1994) The Journal of Biological Chem. 269: 32264-32271; Pang et al. (1997) The Journal of Biological Chem. 272: 25623-25627). Another feature of the invention is the discovery that the 2490 gene is essential for seedling growth and development. An advantage of the present invention is that the newly discovered essential gene containing a novel herbicidal mode of action enables one skilled in the art to easily and rapidly identify novel herbicides.

A further feature of the invention is the identification of a gene in *Arabidopsis*, herein referred to as the 3963 gene, which encodes a protein with sequence similarity to a number of DNA repair proteins, including Rad32p from *Schizosaccharomyces pombe* (Genbank accession number Q09683); hMre11 from *Homo sapiens* (Genbank accession number U37359); and Mre11p from *Saccharomyces cerevisiae* (Genbank accession number U60829) (Johzuka and Ogawa (1995) Genetics, 139: 1521-1532; Paull and Gellert (1998) Molecular Cell, 1: 969-979). Another feature of the invention is the discovery that the 3963 gene is essential for seedling growth and development. An advantage of the present invention is that the newly discovered essential gene containing a novel herbicidal mode of action enables one skilled in the art to easily and rapidly identify novel herbicides.

A further feature of the invention is the identification of a gene in *Arabidopsis*, herein referred to as the 4036 gene, which encodes a protein with sequence similarity to 1-deoxy-D-xylulose 5-phosphate reductoisomerase from a number of organisms including *Synechocystis* sp. (SWISS-PROTQ55663), *Bacillus subtilis* (SWISS-PROT O31753), and *Escherichia coli* (SWISS-PROT P45568) (Takahashi et al. (1998) Proc. Natl. Acad. Sci. USA, 95: 9879-9884). An important and unexpected feature of the invention is the discovery that the 4036 gene is essential for seedling growth and development. An advantage of the present invention is that the newly discovered essential gene containing a novel herbicidal mode of action enables one skilled in the art to easily and rapidly identify novel herbicides.

One object of the present invention is to provide an essential gene in plants for assay development for inhibitory compounds with herbicidal activity. Genetic results show that when the 245 gene, the 5283 gene, the 2490 gene, the 3963 gene or the 4036 gene is mutated in *Arabidopsis*, the resulting phenotype is seedling lethal in the homozygous state. This suggests a critical role for the gene product encoded by the mutated gene.

Using T-DNA insertion mutagenesis, the inventors of the present invention have demonstrated that the activity encoded by the *Arabidopsis* 245 gene, the *Arabidopsis* 5283 gene, the *Arabidopsis* 2490 gene, the *Arabidopsis* 3963 gene or the *Arabidopsis* 4036 gene (herein referred to as 245, 5283, 2490, 3963 or 4036 activity) is essential in *Arabidopsis* seedlings. This implies that chemicals that inhibit the function of the protein in plants are likely to have detrimental effects on plants and are potentially good herbicide candidates. The present invention therefore provides methods of using a purified protein encoded by the gene sequences described below to identify inhibitors thereof, which can then be used as herbicides to suppress the growth of undesirable vegetation, e.g. in fields where crops are grown, particularly agronomically important crops such as maize and other cereal crops such as wheat, oats, rye, sorghum, rice, barley, millet, turf and forage grasses, and the like, as well as cotton, sugar cane, sugar beet, oilseed rape, and soybeans.

The present invention discloses a nucleotide sequence derived from *Arabidopsis*, designated the 245 gene. The nucleotide sequence of the cDNA clone is set forth in SEQ ID NO:1, and the corresponding amino acid sequence is set forth in SEQ ID NO:2. The nucleotide sequence of the partial genomic DNA sequence is set forth in **SEQ ID NO:12**. The present invention also includes nucleotide sequences substantially similar to those set



forth in SEQ ID NO:1. The present invention also encompasses plant proteins whose amino acid sequence are substantially similar to the amino acid sequences set forth in SEQ ID NO:2. Such proteins can be used in a screening assay to identify inhibitors that are potential herbicides.

The present invention further discloses a nucleotide sequence derived from *Arabidopsis*, designated the 5283 gene. The nucleotide sequence of the cDNA clone is set forth in **SEQ ID NO:3**, and the corresponding amino acid sequence is set forth in **SEQ ID NO:4**. The nucleotide sequence of the genomic DNA sequence is set forth in **SEQ ID NO:14**. The present invention also includes nucleotide sequences substantially similar to those set forth in **SEQ ID NO:3**. The present invention also encompasses plant proteins whose amino acid sequence are substantially similar to the amino acid sequences set forth in **SEQ ID NO:4**. Such proteins can be used in a screening assay to identify inhibitors that are potential herbicides.

The present invention further discloses a nucleotide sequence derived from *Arabidopsis*, designated the 2490 gene. The nucleotide sequence of the cDNA clone is set forth in **SEQ ID NO:5**, and the corresponding amino acid sequence is set forth in **SEQ ID NO:6**. The nucleotide sequence of the genomic DNA sequence is set forth in **SEQ ID NO:19**. The present invention also includes nucleotide sequences substantially similar to those set forth in **SEQ ID NO:5**. The present invention also encompasses plant proteins whose amino acid sequence are substantially similar to the amino acid sequences set forth in **SEQ ID NO:6**. Such proteins can be used in a screening assay to identify inhibitors that are potential herbicides.

The present invention further discloses a nucleotide sequence derived from *Arabidopsis*, designated the 3963 gene. The nucleotide sequence of the cDNA clone is set forth in **SEQ ID NO:7**, and the corresponding amino acid sequence is set forth in **SEQ ID NO:8**. The nucleotide sequence of the genomic DNA sequence is set forth in **SEQ ID NO:24**, which contains genomic DNA sequences from both the portion of the MDK4 clone annotated as MDK4.6 and added sequences on the 3' end based on the inventors' reported cDNA clone. The present invention also includes nucleotide sequences substantially similar to those set forth in **SEQ ID NO:7**. The present invention also encompasses plant proteins whose amino acid sequence are substantially similar to the amino acid sequences set forth in **SEQ**

**ID NO:8.** Such proteins can be used in a screening assay to identify inhibitors that are potential herbicides.

The present invention further discloses a nucleotide sequence derived from *Arabidopsis*, designated the 4036 gene. The nucleotide sequence of the cDNA clone is set forth in **SEQ ID NO:9**, and the corresponding amino acid sequence is set forth in **SEQ ID NO:10**. The nucleotide sequence of the genomic DNA sequence is set forth in **SEQ ID NO:27**. Thirteen nucleotide differences are observed by comparing the cDNA clone, derived from cv. Landsberg, and the genomic sequence, derived from cv. Columbia; and Table 1, below, further identifies these differences. **SEQ ID NO:28** is the same as **SEQ ID NO:9**, but with these thirteen nucleotide differences. The corresponding amino acid sequence of **SEQ ID NO:28** is set forth in **SEQ ID NO:29**. The present invention also includes nucleotide sequences substantially similar to those set forth in **SEQ ID NO:9**. The present invention also encompasses plant proteins whose amino acid sequence are substantially similar to the amino acid sequences set forth in **SEQ ID NO:10** and **SEQ ID NO:29**. Such proteins can be used in a screening assay to identify inhibitors that are potential herbicides.

In a preferred embodiment, the present invention relates to a method for identifying chemicals having the ability to inhibit 245, 5283, 2490, 3963 or 4036 activity in plants preferably comprising the steps of: a) obtaining transgenic plants, plant tissue, plant seeds or plant cells, preferably stably transformed, comprising a non-native nucleotide sequence encoding an enzyme having 245, 5283, 2490, 3963 or 4036 activity and capable of overexpressing an enzymatically active 245, 5283, 2490, 3963 or 4036 gene product (either full length or truncated but still active); b) applying a chemical to the transgenic plants, plant cells, tissues or parts and to the isogenic non-transformed plants, plant cells, tissues or parts; c) determining the growth or viability of the transgenic and non-transformed plants, plant cells, tissues after application of the chemical; d) comparing the growth or viability of the transgenic and non-transformed plants, plant cells, tissues after application of the chemical; and e) selecting chemicals that suppress the viability or growth of the non-transgenic plants, plant cells, tissues or parts, without significantly suppressing the growth of the viability or growth of the isogenic transgenic plants, plant cells, tissues or parts. In a preferred embodiment, the enzyme having 245, 5283, 2490, 3963 or 4036 activity is encoded by a nucleotide sequence derived from a plant, preferably *Arabidopsis thaliana*, desirably identical or substantially similar to the nucleotide sequence set forth in **SEQ ID**

NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, respectively. In another embodiment, the enzyme having 245, 5283, 2490, 3963 or 4036 activity is encoded by a nucleotide sequence capable of encoding the amino acid sequence of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10 respectively. In yet another embodiment, the enzyme having 245, 5283, 2490, 3963 or 4036 activity has an amino acid sequence identical or substantially similar to the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10 respectively.

The present invention further embodies plants, plant tissues, plant seeds, and plant cells that have modified 245, 5283, 2490, 3963 or 4036 activity and that are therefore tolerant to inhibition by a herbicide at levels normally inhibitory to naturally occurring 245, 5283, 2490, 3963 or 4036 activity. Herbicide tolerant plants encompassed by the invention include those that would otherwise be potential targets for normally inhibiting herbicides, particularly the agronomically important crops mentioned above. According to this embodiment, plants, plant tissue, plant seeds, or plant cells are transformed, preferably stably transformed, with a recombinant DNA molecule comprising a suitable promoter functional in plants operatively linked to a nucleotide coding sequence that encodes a modified 245, 5283, 2490, 3963 or 4036 gene that is tolerant to inhibition by a herbicide at a concentration that would normally inhibit the activity of wild-type, unmodified 245, 5283, 2490, 3963 or 4036 gene product. Modified 245, 5283, 2490, 3963 or 4036 activity may also be conferred upon a plant by increasing expression of wild-type herbicide-sensitive 245, 5283, 2490, 3963 or 4036 protein by providing multiple copies of wild-type 245, 5283, 2490, 3963 or 4036 genes to the plant or by overexpression of wild-type 245, 5283, 2490, 3963 or 4036 genes under control of a stronger-than-wild-type promoter. The transgenic plants, plant tissue, plant seeds, or plant cells thus created are then selected by conventional selection techniques, whereby herbicide tolerant lines are isolated, characterized, and developed. Alternately, random or site-specific mutagenesis may be used to generate herbicide tolerant lines.

Therefore, the present invention provides a plant, plant cell, plant seed, or plant tissue transformed with a DNA molecule comprising a nucleotide sequence isolated from a plant that encodes an enzyme having 245, 5283, 2490, 3963 or 4036 activity, wherein the DNA expresses the 245, 5283, 2490, 3963 or 4036 activity and wherein the DNA molecule confers upon the plant, plant cell, plant seed, or plant tissue tolerance to a herbicide in amounts that normally inhibits naturally occurring 245, 5283, 2490, 3963 or 4036 activity. According to one example of this embodiment, the enzyme having 245, 5283, 2490, 3963

or 4036 activity is encoded by a nucleotide sequence identical or substantially similar to the nucleotide sequence set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, respectively, or has an amino acid sequence identical or substantially similar to the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, or SEQ ID NO:10, respectively.

The invention also provides a method for suppressing the growth of a plant comprising the step of applying to the plant a chemical that inhibits the naturally occurring 245, 5283, 2490, 3963 or 4036 activity in the plant. In a related aspect, the present invention is directed to a method for selectively suppressing the growth of undesired vegetation in a field containing a crop of planted crop seeds or plants, comprising the steps of: (a) optionally planting herbicide tolerant crops or crop seeds, which are plants or plant seeds that are tolerant to a herbicide that inhibits the naturally occurring 245, 5283, 2490, 3963 or 4036 activity; and (b) applying to the herbicide tolerant crops or crop seeds and the undesired vegetation in the field a herbicide in amounts that inhibit naturally occurring 245, 5283, 2490, 3963 or 4036 activity, wherein the herbicide suppresses the growth of the weeds without significantly suppressing the growth of the crops.

Other objects and advantages of the present invention will become apparent to those skilled in the art from a study of the following description of the invention and non-limiting examples.

As shown in the examples below, the identification of a novel gene structure, as well as the essentiality of the 245 gene, 5283 gene, 2490 gene, 3963 gene or 4036 gene for normal plant growth and development, have been demonstrated for the first time in *Arabidopsis* using T-DNA insertion mutagenesis. Having established the essentiality of 245, 5283, 2490, 3963 or 4036 function in plants and having identified the genes encoding these essential activities, the inventors thereby provide an important and sought after tool for new herbicide development.

*Arabidopsis* insertional mutant lines segregating for seedling lethal mutations are identified as a first step in the identification of essential proteins. Starting with T2 seeds collected from single T1 plants containing T-DNA insertions in their genomes, those lines segregating homozygous seedling lethal seedlings are identified. These lines are found by placing seeds onto minimal plant growth media, which contains the fungicides benomyl and maxim, and screening for inviable seedlings after 7 and 14 days in the light at room

temperature. Inviabile phenotypes include altered pigmentation or altered morphology. These phenotypes are observed either on plates directly or in soil following transplantation of seedlings.

When a line is identified as segregating a seedling lethal, it is determined if the resistance marker in the T-DNA co-segregates with the lethality (Errampalli et al. (1991) *The Plant Cell*, 3:149-157). Co-segregation analysis is done by placing the seeds on media containing the selective agent and scoring the seedlings for resistance or sensitivity to the agent. Examples of selective agents used are hygromycin or phosphinothricin. About 35 resistant seedlings are transplanted to soil and their progeny are examined for the segregation of the seedling lethal. In the case in which the T-DNA insertion disrupts an essential gene, there is co-segregation of the resistance phenotype and the seedling lethal phenotype in every plant. Therefore, in such a case, all resistant plants segregate seedling lethals in the next generation; this result indicates that each of the resistant plants is heterozygous for the DNA causing both phenotypes.

For those lines showing co-segregation of the T-DNA resistance marker and the seedling lethal phenotype, Southern analysis is performed as an initial step in the characterization of the molecular nature of each insertion. Southern analyses are done with genomic DNA isolated from heterozygotes and using probes capable of hybridizing with the T-DNA vector DNA. Using the results of the Southern analysis, appropriate restriction enzymes are chosen to perform plasmid rescue in order to molecularly clone *Arabidopsis* genomic DNA flanking one or both sides of the T-DNA insertion. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of non-T-DNA vector sequences. When such sequences are found, they are used to search DNA and protein databases using the BLAST and BLAST2 programs (Altschul et al. (1990) *J Mol. Biol.* 215: 403-410; Altschul et al (1997) *Nucleic Acid Res.* 25:3389-3402). Additional genomic and cDNA sequences for each gene are identified by standard molecular biology procedures.

The *Arabidopsis* 245 gene was identified by isolating DNA flanking the T-DNA border from the tagged seedling-lethal line #245. A region of the *Arabidopsis* DNA, flanking the T-DNA border, is 99% identical to the genomic survey sequence F17K7TR (accession # B24357). The inventors are the first to demonstrate that the 245 gene product is essential for normal growth and development in plants, as well as defining the function of the 245 gene product

through protein homology. The present invention discloses the cDNA nucleotide sequence of the *Arabidopsis* 245 gene as well as the amino acid sequence of the *Arabidopsis* 245 protein. The nucleotide sequence corresponding to the cDNA clone is set forth in SEQ ID NO:1, and the amino acid sequence encoding the protein is set forth in SEQ ID NO:2. The UTR sequence found 3' to SEQ ID NO:1 is set forth in SEQ ID NO:13. The nucleotide sequence corresponding to the partial genomic DNA is set forth in SEQ ID NO:12. The present invention also encompasses an isolated amino acid sequence derived from a plant, wherein said amino acid sequence is identical or substantially similar to the amino acid sequence encoded by the nucleotide sequence set forth in SEQ ID NO: 1, wherein said amino acid sequence has 245 activity. Using BLAST and BLAST2 programs with the default settings, the sequence of the 245 gene shows similarity to peptide release factor 2 from numerous prokaryotic species. Notable species similarities include: *Escherichia coli* (RF-2) [Swiss-Prot accession #P07012]; *Salmonella typhimurium* (RF-2 Salty)[Swiss-Prot accession # P28353]; and *Mycobacterium tuberculosis* (RF-2: prfB)[Swiss-Prot accession #O05782]. Using GAP analysis of the following protein sequences with the 245 protein results in the following sequence identities with the 245 protein: *Escherichia coli* (RF-2) [Swiss-Prot accession #P07012]( 27.2% identity); *Salmonella typhimurium* (RF-2 Salty)[Swiss-Prot accession # P28353] (24.6% identity); and *Mycobacterium tuberculosis* (RF-2: prfB)[Swiss-Prot accession #O05782] (27.2% identity). In addition, *Synechocystis* (GenPept accession #BAA18577) (31.5% identity); and P1 clone MAB16, chromosome 5 of *Arabidopsis thaliana* (Accession #AB018112NID) (46.2% identity).

The *Arabidopsis* 5283 gene was identified by isolating DNA flanking the T-DNA border from the tagged seedling-lethal line #5283. A region of the *Arabidopsis* DNA, flanking the T-DNA border is identical to an internal region of a sequenced BAC of *Arabidopsis* (BAC T13D8, chromosome 1). This BAC clone contains 116,177 bp of sequence, of which a very small portion corresponds to the genomic region that contains the 5283 gene. Notwithstanding the BAC information, the inventors are the first to demonstrate that the 5283 gene product is essential for normal growth and development in plants, as well as defining the function of the 5283 gene product through protein homology. The present invention discloses the cDNA nucleotide sequence of the *Arabidopsis* 5283 gene as well as the amino acid sequence of the *Arabidopsis* 5283 protein. The nucleotide sequence corresponding to the cDNA clone is set forth in **SEQ ID NO:3**, and the amino acid sequence encoding the protein

is set forth in **SEQ ID NO:4**. The nucleotide sequence corresponding to the genomic DNA is set forth in **SEQ ID NO: 14**. The nucleotide sequence corresponding to the 5' UTR from the cDNA sequence is set forth in **SEQ ID NO:17**, and the nucleotide sequence corresponding to the 3'UTR from the cDNA sequence is set forth in **SEQ ID NO:18**. The present invention also encompasses an isolated amino acid sequence derived from a plant, wherein said amino acid sequence is identical or substantially similar to the amino acid sequence encoded by the nucleotide sequence set forth in **SEQ ID NO: 3**, wherein said amino acid sequence has 5283 activity. Using BLAST and BLAST2 programs with the default settings, the sequence of the 5283 protein shows similarity to SPBC119.13c from *S. pombe* [GENPEPT accession # CAA17928]; SAR DNA-binding proteins from plants [SARBP-1; Genbank accession # AF061962 and SARBP-2: Genbank accession # AF061963]; and prp31 and SIK1p (Nop56) from *S. cerevisiae* [PRP31: Swiss Prot accession # Q12460]. Using GAP analysis of the following protein sequences with the 5283 protein results in the following sequence identities with the 5283 protein: SPBC119.13c from *S. pombe* [GENPEPT accession # CAA17928] (40.5% identity); SAR DNA-binding proteins from plants [SARBP-1; Genbank accession # AF061962 (23.5% identity), and SARBP-2: Genbank accession # AF061963] (24.2% identity); and prp31 and SIK1p (Nop56) from *S. cerevisiae* [PRP31: Swiss Prot accession # Q12460] (24.1% identity). In addition, *Arabidopsis thaliana* (GENPEPT accession # AAC18800) results in 73.8% identity with the 5283 protein.

The *Arabidopsis* 2490 gene is identified by isolating DNA flanking the T-DNA border from the tagged seedling-lethal line #2490. *Arabidopsis* DNA flanking the T-DNA border is identical to an internal region of a sequenced P1 clone of *Arabidopsis* (P1 MTG13, chromosome 5). This P1 clone contains 50,641 bp of sequence, of which a small portion corresponds to the genomic region that contains the 2490 gene. The sequence of a 2490 cDNA containing the entire coding sequence for the 2490 protein is obtained by determining the sequence of the 144K24 EST clone (obtained from Michigan State University). Notwithstanding the BAC and EST sequence information, the inventors are the first to establish definitively the entire gene sequence, and to demonstrate that the 2490 gene product is essential for normal growth and development in plants, as well as defining the function of the 2490 gene product through protein homology. The present invention discloses the cDNA nucleotide sequence of the *Arabidopsis* 2490 gene as well as the

amino acid sequence of the *Arabidopsis* 2490 protein. The nucleotide sequence corresponding to the cDNA clone is set forth in **SEQ ID NO:5**, and the amino acid sequence encoding the protein is set forth in **SEQ ID NO:6**. The UTR sequence found 5' to **SEQ ID NO:5** is set forth in **SEQ ID NO:20**, and the UTR sequence found 3' to **SEQ ID NO:5** is set forth in **SEQ ID NO:21**. The nucleotide sequence corresponding to the genomic DNA is set forth in **SEQ ID NO:19**. The present invention also encompasses an isolated amino acid sequence derived from a plant, wherein said amino acid sequence is identical or substantially similar to the amino acid sequence encoded by the nucleotide sequence set forth in **SEQ ID NO: 5**, wherein said amino acid sequence has 2490 activity. Using BLAST and BLAST2 programs with the default settings, the sequence of the 2490 protein shows similarity to the Toc36 (bce42B) chloroplast envelope protein from *Brassica napus* (Ko et al. (1995) The Journal of Biological Chem. 270: 28601-28608; Wu et al. (1994) The Journal of Biological Chem. 269: 32264-32271; Pang et al. (1997) The Journal of Biological Chem. 272: 25623-25627). Using GAP analysis of the 2490 protein and the Toc36 (bce42B) chloroplast envelope protein from *Brassica napus* (Genbank accession #X79091) results in 81.7% identity with the 2490 protein.

The *Arabidopsis* 3963 gene is identified by isolating DNA flanking the T-DNA border from the tagged seedling-lethal line #3963. A region of the *Arabidopsis* DNA flanking the T-DNA border is 100% identical to the genomic sequence for P1 clone MDK4 on chromosome 5 (Genbank accession number AB010695). The inventors are the first to demonstrate that the 3963 gene product is essential for normal growth and development in plants, as well as defining the function of the 3963 gene product through protein homology. The present invention discloses the cDNA nucleotide sequence of the *Arabidopsis* 3963 gene as well as the amino acid sequence of the *Arabidopsis* 3963 protein. The nucleotide sequence corresponding to the cDNA clone is set forth in **SEQ ID NO:7**, and the amino acid sequence encoding the protein is set forth in **SEQ ID NO:8**. The UTR sequence found 3' to **SEQ ID NO:7** is set forth in **SEQ ID NO:26**. The nucleotide sequence corresponding to the genomic DNA is set forth in **SEQ ID NO:24**. The present invention also encompasses an isolated amino acid sequence derived from a plant, wherein said amino acid sequence is identical or substantially similar to the amino acid sequence encoded by the nucleotide sequence set forth in **SEQ ID NO:7**, wherein said amino acid sequence has 3963 activity. Using BLAST and BLAST2 programs with the default settings, the sequence of the 3963 gene shows



similarity to a number of DNA repair proteins, including Rad32p from *Schizosaccharomyces pombe* (Genbank accession number Q09683); hMre11 from *Homo sapiens* (Genbank accession number U37359); and Mre11p from *Saccharomyces cerevisiae* (Genbank accession number U60829). Using GAP analysis of the following protein sequences with the 3963 protein results in the following sequence identities with the 3963 protein: Rad32p from *Schizosaccharomyces pombe* (Genbank accession number Q09683) (37.5% identity); hMre11 from *Homo sapiens* (Genbank accession number U37359) (39.4% identity); and Mre11p from *Saccharomyces cerevisiae* (Genbank accession number U60829) (34.7% identity).

The *Arabidopsis* 4036 gene is identified by isolating DNA flanking the T-DNA border from the tagged seedling-lethal line #4036. A region of the *Arabidopsis* DNA flanking the T-DNA border is 100% identical to the published genomic sequence for P1 clone MQB2, from chromosome 5 of *Arabidopsis* (Genbank accession # AB009053). The Inventors are the first to demonstrate that the 4036 gene product is essential for normal growth and development in plants, as well as defining the function of the 4036 gene through protein homology. The present invention discloses the cDNA coding nucleotide sequence of the *Arabidopsis* 4036 gene as well as the amino acid sequence of the *Arabidopsis* 4036 protein. The nucleotide sequences corresponding to the cDNA of cv. Landsberg and that of two cultivars are set forth in **SEQ ID NO:9** and **SEQ ID NO:28**, respectively. The corresponding amino acid sequences encoding the proteins are set forth in **SEQ ID NO:10** and **SEQ ID NO:29**. The nucleotide sequence corresponding to the genomic DNA is set forth in **SEQ ID NO:27**. Thirteen nucleotide differences are observed by comparing the cDNA clone, derived from cv. Landsberg, and the genomic sequence, derived from cv. Columbia, and these variations are listed below in Table 1.

Table 1. Nucleotide Differences Observed Between the 4036 cDNA Clone, from cv. Landsberg, and the 4036 Genomic Sequence, from cv. Columbia

Nucleotide #*	cv. Landsberg	cv. Columbia	Codon containing nucleotide difference
			(amino acid residue in cv. Landsberg and amino acid residue in cv. Columbia)**

115	G	A	GAT to AAT (Asp to Asn)
207	T	C	GTT to GTC (Val to Val)
273	C	T	TCC to TCT (Ser to Ser)
276	C	T	ATC to ATT (Ile to Ile)
321	T	C	TTT to TTC (Phe to Phe)
393	G	A	GCG to GCA (Ala to Ala)
485	T	A	CTA to CAA (Leu to Gln)
464	C	T	CCC to CTC (Pro to Leu)
559	A	C	AAG to CAG (Lys to Gln)
963	T	G	CCT to CCG (Pro to Pro)
1101	T	A	CCT to CCA (Pro to Pro)
1254	T	C	TTT to TTC (Phe to Phe)
1393	G	A	GAT to AAT (Asp to Asn)

**\*SEQ ID NO:9** used as a reference for nucleotide numbering

**\*\*Amino acid residues:** Ala (alanine); Asn (asparagine); Asp (aspartic acid); Gln (glutamine); Ile (isoleucine); Leu (leucine); Lys (lysine); Phe (phenylalanine); Pro (proline); Ser (serine); and Val (valine)

The present invention also encompasses an isolated amino acid sequence derived from a plant, wherein said amino acid sequence is identical or substantially similar to the amino acid sequence encoded by the nucleotide sequence set forth in **SEQ ID NO:9**, wherein said amino acid sequence has 4036 activity. Using BLAST and BLAST2 programs with the default settings, the sequence of the 4036 gene shows similarity to 1-deoxy-D-xylulose 5-phosphate reductoisomerase from a number of organisms including *Synechocystis* sp. (SWISS-PROTQ55663), *Bacillus subtilis* (SWISS-PROT O31753); and *Escherichia coli* (SWISS-PROT P45568) (Takahashi et al. (1998) Proc. Natl. Acad. Sci. USA, 95: 9879-9884). Using GAP analysis of the following protein sequences with the 4036 protein results in the following sequence identities with the 4036 protein: 1-deoxy-D-xylulose 5-phosphate reductoisomerase from *Synechocystis* sp. (SWISS-PROTQ55663) (66.1% identity); *Bacillus subtilis* (SWISS-PROT O31753) (45.4% identity); and *Escherichia coli* (SWISS-PROT P45568) (44.6% identity) (Takahashi et al. (1998) Proc. Natl. Acad. Sci. USA, 95: 9879-9884).

For recombinant production of 245, 5283, 2490, 3963 or 4036 activity in a host organism, a nucleotide sequence encoding a protein having 245, 5283, 2490, 3963 or 4036 activity is inserted into an expression cassette designed for the chosen host and introduced into the host where it is recombinantly produced. For example, SEQ ID NO:1 or SEQ ID NO:1 associated with **SEQ ID NO:13** as a 3' UTR, nucleotide sequences substantially similar to SEQ ID NO:1, or homologs of the 245 coding sequence can be used for the recombinant production of a protein having 245 activity. The choice of specific regulatory sequences such as promoter, signal sequence, 5' and 3' untranslated sequences, and enhancer appropriate for the chosen host is within the level of skill of the routineer in the art. The resultant molecule, containing the individual elements operably linked in proper reading frame, may be inserted into a vector capable of being transformed into the host cell. Suitable expression vectors and methods for recombinant production of proteins are well known for host organisms such as *E. coli*, yeast, and insect cells (see, e.g., Luckow and Summers, *Bio/Technol.* 6: 47 (1988), and baculovirus expression vectors, e.g., those derived from the genome of *Autographica californica* nuclear polyhedrosis virus (AcMNPV). A preferred baculovirus/insect system is pAChLT (Pharmingen, San Diego, CA) used to transfect *Spodoptera frugiperda* Sf9 cells (ATCC) in the presence of linear *Autographa californica* baculovirus DNA (Pharmigen, San Diego, CA). The resulting virus is used to infect HighFive *Tricoplusia ni* cells (Invitrogen, La Jolla, CA). In a similar fashion, recombinant production of 5283, 2490, 3963, or 4036 activity is obtained.

In a preferred embodiment, the nucleotide sequence encoding a protein having 245, 5283, 2490, 3963 or 4036 activity is derived from an eukaryote, such as a mammal, a fly or a yeast, but is preferably derived from a plant. In a further preferred embodiment, the nucleotide sequence is identical or substantially similar to the nucleotide sequence set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9 respectively or encodes a protein having 245, 5283, 2490, 3963 or 4036 activity, respectively, whose amino acid sequence is identical or substantially similar to the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10 respectively. The nucleotide sequence set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:9 encodes the *Arabidopsis* 245 protein, *Arabidopsis* 5283 protein, *Arabidopsis* 2490 protein, *Arabidopsis* 3963 protein or *Arabidopsis* 4036 protein, whose amino acid sequence is set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID

NO:6, SEQ ID NO:8 or SEQ ID NO:10 respectively. In another preferred embodiment, the nucleotide sequence is derived from a prokaryote, preferably a bacteria, e.g. *E. coli*.

Recombinantly produced protein having 245, 5283, 2490, 3963 or 4036 activity is isolated and purified using a variety of standard techniques. The actual techniques that may be used will vary depending upon the host organism used, whether the protein is designed for secretion, and other such factors familiar to the skilled artisan (see, e.g. chapter 16 of Ausubel, F. *et al.*, "Current Protocols in Molecular Biology", pub. by John Wiley & Sons, Inc. (1994).

Recombinantly produced proteins having 245, 5283, 2490, 3963 or 4036 activity are useful for a variety of purposes. For example, they can be used in *in vitro* assays to screen known herbicidal chemicals whose target has not been identified to determine if they inhibit 245, 5283, 2490, 3963 or 4036 activity. Such *in vitro* assays may also be used as more general screens to identify chemicals that inhibit such enzymatic activity and that are therefore novel herbicide candidates. Alternatively, recombinantly produced proteins having 245, 5283, 2490, 3963 or 4036 activity may be used to elucidate the complex structure of these molecules and to further characterize their association with known inhibitors in order to rationally design new inhibitory herbicides as well as herbicide tolerant forms of the enzymes.

*In Vitro* Inhibitor Assays: Discovery of Small Molecule Ligand that Interacts with the Gene Product of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9 respectively

Once a protein has been identified as a potential herbicide target, the next step is to develop an assay that allows screening a large number of chemicals to determine which ones interact with the protein. Although it is straightforward to develop assays for proteins of known function, developing assays with proteins of unknown functions is more difficult.

This difficulty can be overcome by using technologies that can detect interactions between a protein and a compound without knowing the biological function of the protein. A short description of three methods is presented, including fluorescence correlation spectroscopy, surface-enhanced laser desorption/ionization, and biacore technologies.

Fluorescence Correlation Spectroscopy (FCS) theory was developed in 1972 but it is only in recent years that the technology to perform FCS became available (Madge et al. (1972) Phys. Rev. Lett., 29: 705-708; Maiti et al. (1997) Proc. Natl. Acad. Sci. USA, 94: 11753-11757). FCS measures the average diffusion rate of a fluorescent molecule within a small sample volume. The sample size can be as low as  $10^3$  fluorescent molecules and the sample volume as low as the cytoplasm of a single bacterium. The diffusion rate is a function of the mass of the molecule and decreases as the mass increases. FCS can therefore be applied to protein-ligand interaction analysis by measuring the change in mass and therefore in diffusion rate of a molecule upon binding. In a typical experiment, the target to be analyzed is expressed as a recombinant protein with a sequence tag, such as a poly-histidine sequence, inserted at the N or C-terminus. The expression takes place in *E. coli*, yeast or insect cells. The protein is purified by chromatography. For example, the poly-histidine tag can be used to bind the expressed protein to a metal chelate column such as  $Ni^{2+}$  chelated on iminodiacetic acid agarose. The protein is then labeled with a fluorescent tag such as carboxytetramethylrhodamine or BODIPY® (Molecular Probes, Eugene, OR). The protein is then exposed in solution to the potential ligand, and its diffusion rate is determined by FCS using instrumentation available from Carl Zeiss, Inc. (Thornwood, NY). Ligand binding is determined by changes in the diffusion rate of the protein.

Surface-Enhanced Laser Desorption/Ionization (SELDI) was invented by Hutchens and Yip during the late 1980's (Hutchens and Yip (1993) Rapid Commun. Mass Spectrom. 7: 576-580). When coupled to a time-of-flight mass spectrometer (TOF), SELDI provides a mean to rapidly analyze molecules retained on a chip. It can be applied to ligand-protein interaction analysis by covalently binding the target protein on the chip and analyze by MS the small molecules that bind to this protein (Worrall et al. (1998) Anal. Biochem. 70: 750-756). In a typical experiment, the target to be analyzed is expressed as described for FCS. The purified protein is then used in the assay without further preparation. It is bound to the SELDI chip either by utilizing the poly-histidine tag or by other interaction such as ion exchange or hydrophobic interaction. The chip thus prepared is then exposed to the potential ligand via, for example, a delivery system capable to pipet the ligands in a sequential manner (autosampler). The chip is then submitted to washes of increasing stringency, for example a series of washes with buffer solutions containing an increasing ionic strength. After each wash, the bound material is analyzed by submitting the chip to SELDI-TOF. Ligands that specifically bind the target will be identified by the stringency of the wash needed to elute them.

Biacore relies on changes in the refractive index at the surface layer upon binding of a ligand to a protein immobilized on the layer. In this system, a collection of small ligands is injected sequentially in a 2-5 ul cell with the immobilized protein. Binding is detected by surface plasmon resonance (SPR) by recording laser light refracting from the surface. In general, the refractive index change for a given change of mass concentration at the surface layer, is practically the same for all proteins and peptides, allowing a single method to be applicable for any protein (Liedberg et al. (1983) *Sensors Actuators* 4: 299-304; Malmquist (1993) *Nature*, 361: 186-187). In a typical experiment, the target to be analyzed is expressed as described for FCS. The purified protein is then used in the assay without further preparation. It is bound to the Biacore chip either by utilizing the poly-histidine tag or by other interaction such as ion exchange or hydrophobic interaction. The chip thus prepared is then exposed to the potential ligand via the delivery system incorporated in the instruments sold by Biacore (Uppsala, Sweden) to pipet the ligands in a sequential manner (autosampler). The SPR signal on the chip is recorded and changes in the refractive index indicate an interaction between the immobilized target and the ligand. Analysis of the signal kinetics on rate and off rate allows the discrimination between non-specific and specific interaction.

Also, an assay for small molecule ligands that interact with a polypeptide is an inhibitor assay. For example, such an inhibitor assay useful for identifying inhibitors of essential plant genes, such as 245, 5283, 2490, 3963, or 4036 genes, comprises the steps of:

- a) reacting a plant 245, 5283, 2490, 3963, or 4036 protein and a substrate thereof in the presence of a suspected inhibitor of the protein's function;
- b) comparing the rate of enzymatic activity in the presence of the suspected inhibitor to the rate of enzymatic activity under the same conditions in the absence of the suspected inhibitor; and
- c) determining whether the suspected inhibitor inhibits the 245, 5283, 2490, 3963, or 4036 protein .

For example, the inhibitory effect on plant 245, 5283, 2490, 3963, or 4036 protein may be determined by a reduction or complete inhibition of 245, 5283, 2490, 3963, or 4036 activity in the assay. Such a determination may be made by comparing, in the presence and absence of the candidate inhibitor, the amount of substrate used or intermediate or product made during the reaction.

In one embodiment, a suspected herbicide, for example identified by *in vitro* screening, is applied to plants at various concentrations. The suspected herbicide is preferably sprayed on the plants. After application of the suspected herbicide, its effect on the plants, for example death or suppression of growth, is recorded.

In another embodiment, an *in vivo* screening assay for inhibitors of the 245, 5283, 2490, 3963 or 4036 activity uses transgenic plants, plant tissue, plant seeds or plant cells capable of overexpressing a nucleotide sequence having 245, 5283, 2490, 3963 or 4036 activity, wherein the 245, 5283, 2490, 3963 or 4036 gene product is enzymatically active in the transgenic plants, plant tissue, plant seeds or plant cells. The nucleotide sequence is preferably derived from an eukaryote, such as a yeast, but is preferably derived from a plant. In a further preferred embodiment, the nucleotide sequence is identical or substantially similar to the nucleotide sequence set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, or encodes an enzyme having 245, 5283, 2490, 3963 or 4036 activity, whose amino acid sequence is identical or substantially similar to the amino acid sequence set forth in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:10 respectively. In another preferred embodiment, the nucleotide sequence is derived from a prokaryote, preferably a bacteria, e.g. *E. coli*.

A chemical is then applied to the transgenic plants, plant tissue, plant seeds or plant cells and to the isogenic non-transgenic plants, plant tissue, plant seeds or plant cells, and the growth or viability of the transgenic and non-transformed plants, plant tissue, plant seeds or plant cells are determined after application of the chemical and compared. Compounds capable of inhibiting the growth of the non-transgenic plants, but not affecting the growth of the transgenic plants are selected as specific inhibitors of 245, 5283, 2490, 3963 or 4036 activity.

The present invention is further directed to plants, plant tissue, plant seeds, and plant cells tolerant to herbicides that inhibit the naturally occurring 245, 5283, 2490, 3963 or 4036 activity in these plants, wherein the tolerance is conferred by an altered 245, 5283, 2490, 3963 or 4036 activity respectively. Altered 245, 5283, 2490, 3963 or 4036 activity may be conferred upon a plant according to the invention by increasing expression of wild-type herbicide-sensitive 245, 5283, 2490, 3963 or 4036 gene, for example by providing additional wild-type 245, 5283, 2490, 3963 or 4036 genes and/or by overexpressing the

endogenous 245, 5283, 2490, 3963 or 4036 gene respectively, for example by driving expression with a strong promoter. Altered 245, 5283, 2490, 3963 or 4036 activity also may be accomplished by expressing nucleotide sequences that are substantially similar to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9 respectively or homologs thereof in a plant. Still further altered 245, 5283, 2490, 3963 or 4036 activity is conferred on a plant by expressing modified herbicide-tolerant 245, 5283, 2490, 3963 or 4036 genes respectively in the plant. Combinations of these techniques may also be used. Representative plants include any plants to which these herbicides are applied for their normally intended purpose. Preferred are agronomically important crops such as cotton, soybean, oilseed rape, sugar beet, maize, rice, wheat, barley, oats, rye, sorghum, millet, turf, forage, turf grasses, and the like.

Achieving altered 245 activity or 5283, 2490, 3963 or 4036 activity respectively through increased expression results in a level of 245 activity or 5283, 2490, 3963, 4036 activity respectively in the plant cell at least sufficient to overcome growth inhibition caused by the herbicide when applied in amounts sufficient to inhibit normal growth of control plants. The level of expressed enzyme generally is at least two times, preferably at least five times, and more preferably at least ten times the natively expressed amount. Increased expression may be due to multiple copies of a wild-type 245 gene or 5283, 2490, 3963 or 4036 gene respectively; multiple occurrences of the coding sequence within the gene (*i.e.* gene amplification) or a mutation in the non-coding, regulatory sequence of the endogenous gene in the plant cell. Plants having such altered gene activity can be obtained by direct selection in plants by methods known in the art (see, *e.g.* U.S. Patent No. 5,162,602, and U.S. Patent No. 4,761,373, and references cited therein). These plants also may be obtained by genetic engineering techniques known in the art. Increased expression of a herbicide-sensitive 245 gene or 5283, 2490, 3963 or 4036 gene respectively can also be accomplished by transforming a plant cell with a recombinant or chimeric DNA molecule comprising a promoter capable of driving expression of an associated structural gene in a plant cell operatively linked to a homologous or heterologous structural gene encoding the 245 protein or the 5283, 2490, 3963 or 4036 protein respectively or a homolog thereof. Preferably, the transformation is stable, thereby providing a heritable transgenic trait.

According to this embodiment, plants, plant tissue, plant seeds, or plant cells are stably transformed with a recombinant DNA molecule comprising a suitable promoter



functional in plants operatively linked to a coding sequence encoding a herbicide tolerant form of the 245, 5283, 2490, 3963 or 4036 protein respectively. A herbicide tolerant form of the enzyme has at least one amino acid substitution, addition or deletion that confers tolerance to a herbicide that inhibits the unmodified, naturally occurring form of the enzyme. The transgenic plants, plant tissue, plant seeds, or plant cells thus created are then selected by conventional selection techniques, whereby herbicide tolerant lines are isolated, characterized, and developed. Below are described methods for obtaining genes that encode herbicide tolerant forms of 245, 5283, 2490, 3963 or 4036 protein respectively:

One general strategy involves direct or indirect mutagenesis procedures on microbes. For instance, a genetically manipulatable microbe such as *E. coli* or *S. cerevisiae* may be subjected to random mutagenesis *in vivo* with mutagens such as UV light or ethyl or methyl methane sulfonate. Mutagenesis procedures are described, for example, in Miller, *Experiments in Molecular Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1972); Davis *et al.*, *Advanced Bacterial Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1980); Sherman *et al.*, *Methods in Yeast Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1983); and U.S. Patent No. 4,975,374. The microbe selected for mutagenesis contains a normal, inhibitor-sensitive 245, 5283, 2490, 3963 or 4036 gene respectively and is dependent upon the activity conferred by this gene. The mutagenized cells are grown in the presence of the inhibitor at concentrations that inhibit the unmodified gene. Colonies of the mutagenized microbe that grow better than the unmutagenized microbe in the presence of the inhibitor (i.e. exhibit resistance to the inhibitor) are selected for further analysis. 245, 5283, 2490, 3963 or 4036 genes respectively conferring tolerance to the inhibitor are isolated from these colonies, either by cloning or by PCR amplification, and their sequences are elucidated. Sequences encoding altered gene products are then cloned back into the microbe to confirm their ability to confer inhibitor tolerance.

A method of obtaining mutant herbicide-tolerant alleles of a plant 245, 5283, 2490, 3963 or 4036 gene involves direct selection in plants. For example, the effect of a mutagenized 245, 5283, 2490, 3963 or 4036 gene on the growth inhibition of plants such as *Arabidopsis*, soybean, or maize is determined by plating seeds sterilized by art-recognized methods on plates on a simple minimal salts medium containing increasing concentrations of the inhibitor. Such concentrations are in the range of 0.001, 0.003, 0.01, 0.03, 0.1, 0.3, 1, 3, 10, 30, 110, 300, 1000 and 3000 parts per million (ppm). The lowest dose at which

significant growth inhibition can be reproducibly detected is used for subsequent experiments. Determination of the lowest dose is routine in the art.

Mutagenesis of plant material is utilized to increase the frequency at which resistant alleles occur in the selected population. Mutagenized seed material is derived from a variety of sources, including chemical or physical mutagenesis of seeds, or chemical or physical mutagenesis of pollen (Neuffer, In *Maize for Biological Research* Sheridan, ed. Univ. Press, Grand Forks, ND., pp. 61-64 (1982)), which is then used to fertilize plants and the resulting M<sub>1</sub> mutant seeds collected. Typically for *Arabidopsis*, M<sub>2</sub> seeds (Lehle Seeds, Tucson, AZ), which are progeny seeds of plants grown from seeds mutagenized with chemicals, such as ethyl methane sulfonate, or with physical agents, such as gamma rays or fast neutrons, are plated at densities of up to 10,000 seeds/plate (10 cm diameter) on minimal salts medium containing an appropriate concentration of inhibitor to select for tolerance. Seedlings that continue to grow and remain green 7-21 days after plating are transplanted to soil and grown to maturity and seed set. Progeny of these seeds are tested for tolerance to the herbicide. If the tolerance trait is dominant, plants whose seed segregate 3:1 / resistant:sensitive are presumed to have been heterozygous for the resistance at the M<sub>2</sub> generation. Plants that give rise to all resistant seed are presumed to have been homozygous for the resistance at the M<sub>2</sub> generation. Such mutagenesis on intact seeds and screening of their M<sub>2</sub> progeny seed can also be carried out on other species, for instance soybean (see, e.g. U.S. Pat. No. 5,084,082). Alternatively, mutant seeds to be screened for herbicide tolerance are obtained as a result of fertilization with pollen mutagenized by chemical or physical means.

Confirmation that the genetic basis of the herbicide tolerance is a 245, 5283, 2490, 3963 or 4036 gene respectively is ascertained as exemplified below. First, alleles of the 245 5283, 2490, 3963 or 4036 gene respectively from plants exhibiting resistance to the inhibitor are isolated using PCR with primers based either upon the *Arabidopsis* cDNA coding sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9 respectively or, more preferably, based upon the unaltered 245, 5283, 2490, 3963 or 4036 gene sequence from the plant used to generate tolerant alleles. After sequencing the alleles to determine the presence of mutations in the coding sequence, the alleles are tested for their ability to confer tolerance to the inhibitor on plants into which the putative tolerance-conferring alleles have been transformed. These plants can be either *Arabidopsis* plants or any other plant whose growth is susceptible to the 245, 5283, 2490, 3963 or 4036 inhibitors respectively. Second, the inserted 245, 5283, 2490, 3963 or 4036

genes are mapped relative to known restriction fragment length polymorphisms (RFLPs) (See, for example, Chang *et al. Proc. Natl. Acad. Sci. USA* 85: 6856-6860 (1988); Nam *et al., Plant Cell* 1: 699-705 (1989), cleaved amplified polymorphic sequences (CAPS) (Konieczny and Ausubel (1993) *The Plant Journal*, 4(2): 403-410), or SSLPs (Bell and Ecker (1994) *Genomics*, 19: 137-144). The 245, 5283, 2490, 3963 or 4036 inhibitor tolerance trait respectively is independently mapped using the same markers. When tolerance is due to a mutation in that 245, 5283, 2490, 3963 or 4036 gene respectively, the tolerance trait maps to a position indistinguishable from the position of the 245, 5283, 2490, 3963 or 4036 gene.

Another method of obtaining herbicide-tolerant alleles of a 245, 5283, 2490, 3963 or 4036 gene is by selection in plant cell cultures. Explants of plant tissue, *e.g.* embryos, leaf disks, etc. or actively growing callus or suspension cultures of a plant of interest are grown on medium in the presence of increasing concentrations of the inhibitory herbicide or an analogous inhibitor suitable for use in a laboratory environment. Varying degrees of growth are recorded in different cultures. In certain cultures, fast-growing variant colonies arise that continue to grow even in the presence of normally inhibitory concentrations of inhibitor. The frequency with which such faster-growing variants occur can be increased by treatment with a chemical or physical mutagen before exposing the tissues or cells to the inhibitor. Putative tolerance-conferring alleles of the 245, 5283, 2490, 3963 or 4036 gene respectively are isolated and tested as described in the foregoing paragraphs. Those alleles identified as conferring herbicide tolerance may then be engineered for optimal expression and transformed into the plant. Alternatively, plants can be regenerated from the tissue or cell cultures containing these alleles.

Still another method involves mutagenesis of wild-type, herbicide sensitive plant 245, 5283, 2490, 3963 or 4036 genes respectively in bacteria or yeast, followed by culturing the microbe on medium that contains inhibitory concentrations (i.e. sufficient to cause abnormal growth, inhibit growth or cause cell death) of the inhibitor, and then selecting those colonies that grow normally in the presence of the inhibitor. More specifically, a plant cDNA, such as the *Arabidopsis* cDNA encoding the 245, 5283, 2490, 3963 or 4036 protein respectively, is cloned into a microbe that otherwise lacks the 245 5283, 2490, 3963 or 4036 activity respectively. The transformed microbe is then subjected to *in vivo* mutagenesis or to *in vitro* mutagenesis by any of several chemical or enzymatic methods known in the art, *e.g.* sodium bisulfite (Shortle *et al., Methods Enzymol.* 100:457-468

(1983); methoxylamine (Kadonaga *et al.*, *Nucleic Acids Res.* 13:1733-1745 (1985); oligonucleotide-directed saturation mutagenesis (Hutchinson *et al.*, *Proc. Natl. Acad. Sci. USA*, 83:710-714 (1986); or various polymerase misincorporation strategies (see, e.g. Shortle *et al.*, *Proc. Natl. Acad. Sci. USA*, 79:1588-1592 (1982); Shiraishi *et al.*, *Gene* 64:313-319 (1988); and Leung *et al.*, *Technique* 1:11-15 (1989). Colonies that grow normally in the presence of normally inhibitory concentrations of inhibitor are picked and purified by repeated restreaking. Their plasmids are purified and tested for the ability to confer tolerance to the inhibitor by retransforming them into the microbe lacking 245, 5283, 2490, 3963 or 4036 activity respectively. The DNA sequences of cDNA inserts from plasmids that pass this test are then determined.

Herbicide resistant 245, 5283, 2490, 3963 or 4036 proteins respectively are also obtained using methods involving *in vitro* recombination, also called DNA shuffling. By DNA shuffling, mutations, preferably random mutations, are introduced into nucleotide sequences encoding 245, 5283, 2490, 3963 or 4036 activity respectively. DNA shuffling also leads to the recombination and rearrangement of sequences within a 245, 5283, 2490, 3963 or 4036 gene respectively or to recombination and exchange of sequences between two or more different of 245, 5283, 2490, 3963 or 4036 genes respectively. These methods allow for the production of millions of mutated 245, 5283, 2490, 3963 or 4036 coding sequences respectively. The mutated genes, or shuffled genes, are screened for desirable properties, e.g. improved tolerance to herbicides and for mutations that provide broad spectrum tolerance to the different classes of inhibitor chemistry. Such screens are well within the skills of a routineer in the art.

In a preferred embodiment, a mutagenized 245, 5283, 2490, 3963 or 4036 gene respectively is formed from at least one template 245, 5283, 2490, 3963 or 4036 gene respectively, wherein the template 245 5283, 2490, 3963 or 4036 gene respectively has been cleaved into double-stranded random fragments of a desired size, and comprising the steps of adding to the resultant population of double-stranded random fragments one or more single or double-stranded oligonucleotides, wherein said oligonucleotides comprise an area of identity and an area of heterology to the double-stranded random fragments; denaturing the resultant mixture of double-stranded random fragments and oligonucleotides into single-stranded fragments; incubating the resultant population of single-stranded fragments with a polymerase under conditions which result in the annealing of said single-stranded fragments at said areas of identity to form pairs of annealed fragments, said areas

of identity being sufficient for one member of a pair to prime replication of the other, thereby forming a mutagenized double-stranded polynucleotide; and repeating the second and third steps for at least two further cycles, wherein the resultant mixture in the second step of a further cycle includes the mutagenized double-stranded polynucleotide from the third step of the previous cycle, and the further cycle forms a further mutagenized double-stranded polynucleotide, wherein the mutagenized polynucleotide is a mutated 245, 5283, 2490, 3963 or 4036 gene respectively having enhanced tolerance to a herbicide which inhibits naturally occurring 245, 5283, 2490, 3963 or 4036 activity respectively. In a preferred embodiment, the concentration of a single species of double-stranded random fragment in the population of double-stranded random fragments is less than 1% by weight of the total DNA. In a further preferred embodiment, the template double-stranded polynucleotide comprises at least about 100 species of polynucleotides. In another preferred embodiment, the size of the double-stranded random fragments is from about 5 bp to 5 kb. In a further preferred embodiment, the fourth step of the method comprises repeating the second and the third steps for at least 10 cycles. Such method is described e.g. in Stemmer et al. (1994) Nature 370: 389-391, in US Patent 5,605,793, US Patent 5,811,238 and in Cramer et al. (1998) Nature 391: 288-291, as well as in WO 97/20078, and these references are incorporated herein by reference.

In another preferred embodiment, any combination of two or more different 245 genes are mutagenized *in vitro* by a staggered extension process (StEP), as described e.g. in Zhao et al. (1998) Nature Biotechnology 16: 258-261. The two or more 245 genes are used as template for PCR amplification with the extension cycles of the PCR reaction preferably carried out at a lower temperature than the optimal polymerization temperature of the polymerase. In a similar fashion, the StEP is performed with the 5283, 2490, 3963, or 4036 genes. For example, when a thermostable polymerase with an optimal temperature of approximately 72°C is used, the temperature for the extension reaction is desirably below 72°C, more desirably below 65°C, preferably below 60°C, more preferably the temperature for the extension reaction is 55°C. Additionally, the duration of the extension reaction of the PCR cycles is desirably shorter than usually carried out in the art, more desirably it is less than 30 seconds, preferably it is less than 15 seconds, more preferably the duration of the extension reaction is 5 seconds. Only a short DNA fragment is polymerized in each extension reaction, allowing template switch of the extension products between the starting DNA molecules after each cycle of denaturation and annealing, thereby generating diversity among the extension products. The optimal number of cycles in the PCR reaction depends

on the length of the 245, 5283, 2490, 3963 or 4036 genes respectively to be mutagenized but desirably over 40 cycles, more desirably over 60 cycles, preferably over 80 cycles are used. Optimal extension conditions and the optimal number of PCR cycles for every combination of 245, 5283, 2490, 3963 or 4036 genes respectively are determined as described in using procedures well-known in the art. The other parameters for the PCR reaction are essentially the same as commonly used in the art. The primers for the amplification reaction are preferably designed to anneal to DNA sequences located outside of the 245, 5283, 2490, 3963 or 4036 genes, e.g. to DNA sequences of a vector comprising the 245, 5283, 2490, 3963 or 4036 genes respectively, whereby the different 245, 5283, 2490, 3963 or 4036 genes respectively used in the PCR reaction are preferably comprised in separate vectors. The primers desirably anneal to sequences located less than 500 bp away from 245, 5283, 2490, 3963 or 4036 respectively sequences, preferably less than 200 bp, more preferably less than 120 bp away from the 245, 5283, 2490, 3963 or 4036 sequences respectively. Preferably, the 245, 5283, 2490, 3963 or 4036 sequences respectively are surrounded by restriction sites, which are included in the DNA sequence amplified during the PCR reaction, thereby facilitating the cloning of the amplified products into a suitable vector. In another preferred embodiment, fragments of 245 5283, 2490, 3963 or 4036 genes respectively having cohesive ends are produced as described in WO 98/05765. The cohesive ends are produced by ligating a first oligonucleotide corresponding to a part of a 245, 5283, 2490, 3963 or 4036 gene respectively to a second oligonucleotide not present in the gene or corresponding to a part of the gene not adjoining to the part of the gene corresponding to the first oligonucleotide, wherein the second oligonucleotide contains at least one ribonucleotide. A double-stranded DNA is produced using the first oligonucleotide as template and the second oligonucleotide as primer. The ribonucleotide is cleaved and removed. The nucleotide(s) located 5' to the ribonucleotide is also removed, resulting in double-stranded fragments having cohesive ends. Such fragments are randomly reassembled by ligation to obtain novel combinations of gene sequences.

Any 245, 5283, 2490, 3963 or 4036 gene respectively or any combination of 245 5283, 2490, 3963 or 4036 genes is used for *in vitro* recombination in the context of the present invention, for example, a 245, 5283, 2490, 3963 or 4036 gene respectively derived from a plant, such as, e.g. *Arabidopsis thaliana*, e.g. a 245, 5283, 2490, 3963 or 4036 gene respectively set forth in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9 respectively, or a 245-like, 5283-like, 2490-like, 3963-like or 4036-like gene respectively from *E. coli* (Craigie et al. (1985) Proc Natl Acad Sci, 82: 3616-3620; Craigie

and Caskey (1987) Biochimie, 69: 1031-1041; Ito et al. (1998) Proc Natl Acad Sci, 95: 8165-8169), all of which are incorporated herein by reference. Whole 245, 5283, 2490, 3963 or 4036 genes respectively or portions thereof are used in the context of the present invention. The library of mutated 245, 5283, 2490, 3963 or 4036 genes respectively obtained by the methods described above are cloned into appropriate expression vectors and the resulting vectors are transformed into an appropriate host, for example an algae like *Chlamydomonas*, a yeast or a bacteria. An appropriate host is preferably a host that otherwise lacks 245, 5283, 2490, 3963 or 4036 activity, for example *E. coli*. Host cells transformed with the vectors comprising the library of mutated 245, 5283, 2490, 3963 or 4036 genes respectively are cultured on medium that contains inhibitory concentrations of the inhibitor and those colonies that grow in the presence of the inhibitor are selected. Colonies that grow in the presence of normally inhibitory concentrations of inhibitor are picked and purified by repeated restreaking. Their plasmids are purified and the DNA sequences of cDNA inserts from plasmids that pass this test are then determined.

An assay for identifying a modified 245, 5283, 2490, 3963 or 4036 gene respectively that is tolerant to an inhibitor may be performed in the same manner as the assay to identify inhibitors of the 245, 5283, 2490, 3963 or 4036 activity respectively (Inhibitor Assay, above) with the following modifications: First, a mutant 245, 5283, 2490, 3963 or 4036 protein respectively is substituted in one of the reaction mixtures for the wild-type 245 5283, 2490, 3963 or 4036 protein respectively of the inhibitor assay. Second, an inhibitor of wild-type enzyme is present in both reaction mixtures. Third, mutated activity (activity in the presence of inhibitor and mutated enzyme) and unmutated activity (activity in the presence of inhibitor and wild-type enzyme) are compared to determine whether a significant increase in enzymatic activity is observed in the mutated activity when compared to the unmutated activity. Mutated activity is any measure of activity of the mutated enzyme while in the presence of a suitable substrate and the inhibitor. Unmutated activity is any measure of activity of the wild-type enzyme while in the presence of a suitable substrate and the inhibitor.

In addition to being used to create herbicide-tolerant plants, genes encoding herbicide tolerant 245, 5283, 2490, 3963 or 4036 protein respectively can also be used as selectable markers in plant cell transformation methods. For example, plants, plant tissue, plant seeds, or plant cells transformed with a heterologous DNA sequence can also be transformed with a sequence encoding an altered 245, 5283, 2490, 3963 or 4036 activity respectively capable of being expressed by the plant. The transformed cells are transferred

to medium containing an inhibitor of the enzyme in an amount sufficient to inhibit the growth or survivability of plant cells not expressing the modified coding sequence, wherein only the transformed cells will grow. The method is applicable to any plant cell capable of being transformed with a modified 245, 5283, 2490, 3963 or 4036 gene, and can be used with any heterologous DNA sequence of interest. Expression of the heterologous DNA sequence and the modified gene can be driven by the same promoter functional in plant cells, or by separate promoters.

#### X.

A wild type or herbicide-tolerant form of the 245, 5283, 2490, 3963 or 4036 gene respectively, or homologs thereof, can be incorporated in plant or bacterial cells using conventional recombinant DNA technology. Generally, this involves inserting a DNA molecule encoding the 245, 5283, 2490, 3963 or 4036 gene respectively into an expression system to which the DNA molecule is heterologous (i.e., not normally present) using standard cloning procedures known in the art. The vector contains the necessary elements for the transcription and translation of the inserted protein-coding sequences in a host cell containing the vector. A large number of vector systems known in the art can be used, such as plasmids, bacteriophage viruses and other modified viruses. The components of the expression system may also be modified to increase expression. For example, truncated sequences, nucleotide substitutions, nucleotide optimization or other modifications may be employed. Expression systems known in the art can be used to transform virtually any crop plant cell under suitable conditions. A heterologous DNA sequence comprising a wild-type or herbicide-tolerant form of the 245, 5283, 2490, 3963 or 4036 gene respectively is preferably stably transformed and integrated into the genome of the host cells. In another preferred embodiment, the heterologous DNA sequence comprising a wild-type or herbicide-tolerant form of the 245, 5283, 2490, 3963 or 4036 gene respectively located on a self-replicating vector. Examples of self-replicating vectors are viruses, in particular gemini viruses. Transformed cells can be regenerated into whole plants such that the chosen form of the 245, 5283, 2490, 3963 or 4036 gene respectively confers herbicide tolerance in the transgenic plants.

Gene sequences intended for expression in transgenic plants are first assembled in expression cassettes behind a suitable promoter expressible in plants. The expression



cassettes may also comprise any further sequences required or selected for the expression of the heterologous DNA sequence. Such sequences include, but are not restricted to, transcription terminators, extraneous sequences to enhance expression such as introns, vital sequences, and sequences intended for the targeting of the gene product to specific organelles and cell compartments. These expression cassettes can then be easily transferred to the plant transformation vectors described *infra*. The following is a description of various components of typical expression cassettes.

The selection of the promoter used in expression cassettes will determine the spatial and temporal expression pattern of the heterologous DNA sequence in the plant transformed with this DNA sequence. Selected promoters will express heterologous DNA sequences in specific cell types (such as leaf epidermal cells, mesophyll cells, root cortex cells) or in specific tissues or organs (roots, leaves or flowers, for example) and the selection will reflect the desired location of accumulation of the gene product. Alternatively, the selected promoter may drive expression of the gene under various inducing conditions. Promoters vary in their strength, i.e., ability to promote transcription. Depending upon the host cell system utilized, any one of a number of suitable promoters known in the art can be used. For example, for constitutive expression, the CaMV 35S promoter, the rice actin promoter, or the ubiquitin promoter may be used. For regulatable expression, the chemically inducible PR-1 promoter from tobacco or *Arabidopsis* may be used (*see, e.g.*, U.S. Patent No. 5,689,044).

A variety of transcriptional terminators are available for use in expression cassettes. These are responsible for the termination of transcription beyond the heterologous DNA sequence and its correct polyadenylation. Appropriate transcriptional terminators are those that are known to function in plants and include the CaMV 35S terminator, the *tm1* terminator, the nopaline synthase terminator and the pea *rbcS* E9 terminator. These can be used in both monocotyledonous and dicotyledonous plants.

Numerous sequences have been found to enhance gene expression from within the transcriptional unit and these sequences can be used in conjunction with the genes of this invention to increase their expression in transgenic plants. For example, various intron sequences such as introns of the maize *Adh1* gene have been shown to enhance expression, particularly in monocotyledonous cells. In addition, a number of non-translated

leader sequences derived from viruses are also known to enhance expression, and these are particularly effective in dicotyledonous cells.

The coding sequence of the selected gene optionally is genetically engineered by altering the coding sequence for optimal expression in the crop species of interest. Methods for modifying coding sequences to achieve optimal expression in a particular crop species are well known (see, e.g. Perlak *et al.*, *Proc. Natl. Acad. Sci. USA* 88: 3324 (1991); and Koziel *et al.*, *Bio/technol.* 11: 194 (1993); Fennoy and Bailey-Serres. *Nucl. Acids Res.* 21: 5294-5300 (1993). Methods for modifying coding sequences by taking into account codon usage in plant genes and in higher plants, green algae, and cyanobacteria are well known (see table 4 in: Murray *et al.* *Nucl. Acids Res.* 17: 477-498 (1989); Campbell and Gowri *Plant Physiol.* 92: 1-11(1990).

Various mechanisms for targeting gene products are known to exist in plants and the sequences controlling the functioning of these mechanisms have been characterized in some detail. For example, the targeting of gene products to the chloroplast is controlled by a signal sequence found at the amino terminal end of various proteins which is cleaved during chloroplast import to yield the mature protein (e.g. Comai *et al.* *J. Biol. Chem.* 263: 15104-15109 (1988)). Other gene products are localized to other organelles such as the mitochondrion and the peroxisome (e.g. Unger *et al.* *Plant Molec. Biol.* 13: 411-418 (1989)). The cDNAs encoding these products can also be manipulated to effect the targeting of heterologous products encoded by DNA sequences to these organelles. In addition, sequences have been characterized which cause the targeting of products encoded by DNA sequences to other cell compartments. Amino terminal sequences are responsible for targeting to the ER, the apoplast, and extracellular secretion from aleurone cells (Koehler & Ho, *Plant Cell* 2: 769-783 (1990)). Additionally, amino terminal sequences in conjunction with carboxy terminal sequences are responsible for vacuolar targeting of gene products (Shinshi *et al.* *Plant Molec. Biol.* 14: 357-368 (1990)). By the fusion of the appropriate targeting sequences described above to heterologous DNA sequences of interest it is possible to direct this product to any organelle or cell compartment.

Numerous transformation vectors available for plant transformation are known to those of ordinary skill in the plant transformation arts, and the genes pertinent to this invention can be used in conjunction with any such vectors. The selection of vector will depend upon the preferred transformation technique and the target species for transformation. For certain target species, different antibiotic or herbicide selection markers may be preferred. Selection markers used routinely in transformation include the *nptII* gene, which confers resistance to kanamycin and related antibiotics (Messing & Vierra. Gene 19: 259-268 (1982); Bevan et al., Nature 304:184-187 (1983)), the *bar* gene, which confers resistance to the herbicide phosphinothricin (White et al., Nucl. Acids Res 18: 1062 (1990), Spencer et al. Theor. Appl. Genet 79: 625-631 (1990)), the *hph* gene, which confers resistance to the antibiotic hygromycin (Blochinger & Diggelmann, Mol Cell Biol 4: 2929-2931), and the *dhfr* gene, which confers resistance to methotrexate (Bourouis et al., EMBO J. 2(7): 1099-1104 (1983)), and the EPSPS gene, which confers resistance to glyphosate (U.S. Patent Nos. 4,940,935 and 5,188,642).

Many vectors are available for transformation using *Agrobacterium tumefaciens*. These typically carry at least one T-DNA border sequence and include vectors such as pBIN19 (Bevan, Nucl. Acids Res. (1984)). Typical vectors suitable for *Agrobacterium* transformation include the binary vectors pCIB200 and pCIB2001, as well as the binary vector pCIB10 and hygromycin selection derivatives thereof. (See, for example, U.S. Patent No. 5,639,949).

Transformation without the use of *Agrobacterium tumefaciens* circumvents the requirement for T-DNA sequences in the chosen transformation vector and consequently vectors lacking these sequences can be utilized in addition to vectors such as the ones described above which contain T-DNA sequences. Transformation techniques that do not rely on *Agrobacterium* include transformation via particle bombardment, protoplast uptake (e.g. PEG and electroporation) and microinjection. The choice of vector depends largely on the preferred selection for the species being transformed. Typical vectors suitable for non-

*Agrobacterium* transformation include pCIB3064, pSOG19, and pSOG35. (See, for example, U.S. Patent No. 5,639,949).

Once the coding sequence of interest has been cloned into an expression system, it is transformed into a plant cell. Methods for transformation and regeneration of plants are well known in the art. For example, Ti plasmid vectors have been utilized for the delivery of foreign DNA, as well as direct DNA uptake, liposomes, electroporation, micro-injection, and microprojectiles. In addition, bacteria from the genus *Agrobacterium* can be utilized to transform plant cells.

Transformation techniques for dicotyledons are well known in the art and include *Agrobacterium*-based techniques and techniques that do not require *Agrobacterium*. Non-*Agrobacterium* techniques involve the uptake of exogenous genetic material directly by protoplasts or cells. This can be accomplished by PEG or electroporation mediated uptake, particle bombardment-mediated delivery, or microinjection. In each case the transformed cells are regenerated to whole plants using standard techniques known in the art.

Transformation of most monocotyledon species has now also become routine. Preferred techniques include direct gene transfer into protoplasts using PEG or electroporation techniques, particle bombardment into callus tissue, as well as *Agrobacterium*-mediated transformation.

In another preferred embodiment, a nucleotide sequence encoding a polypeptide having 245, 5283, 2490, 3963, or 4036 activity is directly transformed into the plastid genome. Plastid expression, in which genes are inserted by homologous recombination into the several thousand copies of the circular plastid genome present in each plant cell, takes advantage of the enormous copy number advantage over nuclear-expressed genes to permit expression levels that can readily exceed 10% of the total soluble plant protein. In a preferred embodiment, the nucleotide sequence is inserted into a plastid targeting vector and transformed into the plastid genome of a desired plant host. Plants homoplasmic for plastid genomes containing the nucleotide sequence are obtained, and are preferentially capable of high expression of the nucleotide sequence.

Plastid transformation technology is for example extensively described in U.S. Patent Nos. 5,451,513, 5,545,817, 5,545,818, and 5,877,462 in PCT application no. WO 95/16783 and WO 97/32977, and in McBride *et al.* (1994) Proc. Natl. Acad. Sci. USA 91, 7301-7305, all

incorporated herein by reference in their entirety. The basic technique for plastid transformation involves introducing regions of cloned plastid DNA flanking a selectable marker together with the nucleotide sequence into a suitable target tissue, e.g., using biolistics or protoplast transformation (e.g., calcium chloride or PEG mediated transformation). The 1 to 1.5 kb flanking regions, termed targeting sequences, facilitate homologous recombination with the plastid genome and thus allow the replacement or modification of specific regions of the plastome. Initially, point mutations in the chloroplast 16S rRNA and rps12 genes conferring resistance to spectinomycin and/or streptomycin are utilized as selectable markers for transformation (Svab, Z., Hajdukiewicz, P., and Maliga, P. (1990) *Proc. Natl. Acad. Sci. USA* 87, 8526-8530; Staub, J. M., and Maliga, P. (1992) *Plant Cell* 4, 39-45). The presence of cloning sites between these markers allowed creation of a plastid targeting vector for introduction of foreign genes (Staub, J.M., and Maliga, P. (1993) *EMBO J.* 12, 601-606). Substantial increases in transformation frequency are obtained by replacement of the recessive rRNA or r-protein antibiotic resistance genes with a dominant selectable marker, the bacterial *aadA* gene encoding the spectinomycin-detoxifying enzyme aminoglycoside-3'-adenyltransferase (Svab, Z., and Maliga, P. (1993) *Proc. Natl. Acad. Sci. USA* 90, 913-917). Other selectable markers useful for plastid transformation are known in the art and encompassed within the scope of the invention.

The wild-type or altered form of a 245, 5283, 2490, 3963 or 4036 gene respectively of the present invention can be utilized to confer herbicide tolerance to a wide variety of plant cells, including those of gymnosperms, monocots, and dicots. Although the gene can be inserted into any plant cell falling within these broad classes, it is particularly useful in crop plant cells, such as rice, wheat, barley, rye, corn, potato, carrot, sweet potato, sugar beet, bean, pea, chicory, lettuce, cabbage, cauliflower, broccoli, turnip, radish, spinach, asparagus, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, quince, melon, plum, cherry, peach, nectarine, apricot, strawberry, grape, raspberry, blackberry, pineapple, avocado, papaya, mango, banana, soybean, tobacco, tomato, sorghum and sugarcane.

The high-level expression of a wild-type 245, 5283, 2490, 3963 or 4036 gene respectively and/or the expression of herbicide-tolerant forms of a 245, 5283, 2490, 3963 or 4036 gene respectively conferring herbicide tolerance in plants, in combination with other characteristics important for production and quality, can be incorporated into plant lines through breeding approaches and techniques known in the art.

Where a herbicide tolerant 245, 5283, 2490, 3963 or 4036 gene allele respectively is obtained by direct selection in a crop plant or plant cell culture from which a crop plant can be regenerated, it is moved into commercial varieties using traditional breeding techniques to develop a herbicide tolerant crop without the need for genetically engineering the allele and transforming it into the plant.

The invention will be further described by reference to the following detailed examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified.

## EXAMPLES

Standard recombinant DNA and molecular cloning techniques used here are well known in the art and are described by Sambrook, *et al.*, Molecular Cloning, eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989) and by T.J. Silhavy, M.L. Berman, and L.W. Enquist, Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1984) and by Ausubel, F.M. *et al.*, Current Protocols in Molecular Biology, pub. by Greene Publishing Assoc. and Wiley-Interscience (1987), Reiter, *et al.*, Methods in Arabidopsis Research, World Scientific Press (1992), and Schultz *et al.*, Plant Molecular Biology Manual, Kluwer Academic Publishers (1998). These references describe the standard techniques used for all steps in tagging and cloning genes from T-DNA mutagenized populations of *Arabidopsis*: plant infection and transformation; screening for the identification of seedling mutants; cosegregation analysis; and plasmid rescue.

**Example 1: Sequence Analysis of Tagged Seedling – Lethal Line #245 From the T-DNA Mutagenized Population of *Arabidopsis***

The plasmid rescue technique is used to molecularly clone *Arabidopsis* genomic DNA flanking one or both sides of T-DNA insertions resulting from T-DNA mutagenesis. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of non-T-DNA vector sequences. The plasmids recovered from the plasmid rescue protocol are sequenced using the slp346for primer (SEQ ID NO:11). Primer slp346for provides information on the flanking sequence immediately adjacent to the left T-DNA border. Plasmid rescue is validated by PCR of genomic DNA from a homozygote for the 245 mutation. This PCR experiment uses a primer anchored in the predicted flanking sequence and the slp346for primer (anchored in the T-DNA insertion). Finding a PCR product of the size expected based on the sequence of the plasmid rescued clone confirms a valid rescue. The sequence obtained from primer slp346for is used in a BLASTx search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402.). The BLAST search results show that the recovered plant flanking sequence shows a high level of similarity to numerous prokaryotic

peptide release factor two proteins. The BLAST results indicate that the T-DNA insertion has occurred in the ORF of the first identified plant derived peptide release factor two. A DNA fragment that includes peptide release factor sequence similarity is isolated by amplification of *Arabidopsis* genomic DNA using the polymerase chain reaction. This fragment is used to probe an *Arabidopsis* cDNA library in the  $\lambda$ YES vector (Elledge *et al.* (1991) Proc. Natl. Acad. Sci. 88:1731-1735). Positive phage clones are isolated and characterized using standard molecular biology techniques. The resultant cDNA clones are excised from the phage and the nucleotide sequence is determined. The DNA sequence is shown in SEQ ID NO:1. The deduced amino acid sequence is analyzed using the BLASTx search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402). The BLAST search results show that the recovered 245 cDNA shows sequence similarity to the same set of prokaryotic peptide release factors.

#### Example 2: Sequence Analysis of Tagged Seedling – Lethal Line #5283 From the T-DNA Mutagenized Population of *Arabidopsis*

The plasmid rescue technique is used to molecularly clone *Arabidopsis* genomic DNA flanking one or both sides of T-DNA insertions resulting from T-DNA mutagenesis. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of non-T-DNA vector sequences. The plasmids recovered from the plasmid rescue protocol are sequenced using the slp346for primer (**SEQ ID NO:11**). Primer slp346for provides information on the flanking sequence immediately adjacent to the left T-DNA border. Plasmid rescue is validated by PCR of genomic DNA from a heterozygote for the 5283 mutation. This PCR experiment uses a primer anchored in the predicted flanking sequence and the slp328 primer (**SEQ ID NO:15**) (anchored in the T-DNA insertion). Finding a PCR product of the size expected based on the sequence of the plasmid rescued clone confirms a valid rescue.

The sequence obtained from primer SLP346for is used in a BLASTn search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402.). The BLAST search results show that the



recovered sequence is identical to genomic DNA located in *Arabidopsis* chromosome I, BAC T13D8 (Genbank accession number AC004473). Primer LW60 (**SEQ ID NO:16**), the reverse complement to nucleotides #32,964-32,987 in the BAC T13D8 sequence (5'-aaacgcttacatattcttttcta-3'), is designed and used to determine the sequence downstream of the T-DNA insert; this experiment identifies the junction of the right border. The region of genomic DNA where the T-DNA insertion occurred includes bases #32,879 through #32,885 of the annotated BAC T13D8 sequence, resulting in a six-base deletion. This insertion occurs 90 nucleotides upstream of the sequence annotated on BAC T13D8 as encoding a protein similar to *S. cerevisiae* SIK1P protein (Genbank accession number U20237). A DNA fragment that includes bases #33,025 through bases #34,338 of the BAC T13D8 sequence is isolated by amplification of *Arabidopsis* genomic DNA using the polymerase chain reaction. This fragment is used to probe an *Arabidopsis* cDNA library in the IYES vector (Elledge *et al.* (1991) Proc. Natl. Acad. Sci. 88:1731-1735). Positive phage clones are isolated and characterized using standard molecular biology techniques. The resultant cDNA clones are excised from the phage and the nucleotide sequence is determined. One full-length clone is identified. The deduced amino acid sequence is analyzed using the tBLASTn search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402). The BLAST search results show that the recovered 5283 cDNA sequence is derived from the same genomic sequence located in *Arabidopsis* chromosome I, BAC T13D8. The intron/exon boundaries of the cDNA sequence are the same as those predicted for the *Arabidopsis* SIK1P homolog (Genbank accession number AC004473), with the following exceptions. The initiator codon for the 5283 cDNA is encoded by bases #32975 through #32977, followed immediately by an intron at bases #32978 through #33199.

#### Example 3: Sequence Analysis of Tagged Seedling – Lethal Line #2490 From the T-DNA Mutagenized Population of *Arabidopsis*

The plasmid rescue technique is used to molecularly clone *Arabidopsis* genomic DNA flanking one or both sides of T-DNA insertions resulting from T-DNA mutagenesis. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of

non-T-DNA vector sequences. The plasmids recovered from the plasmid rescue protocol are sequenced using the SLP346for primer (5' GCGGACATCTACATTTTGA 3': **SEQ ID NO:11**). Primer SLP346for provides information on the flanking sequence immediately adjacent to the left T-DNA border. Clones for both ends of the T-DNA insertion are recovered as plasmids containing left T-DNA border. Plasmid rescue is validated by Southern blot analysis comparing genomic DNA from a plant heterozygous for the 2490 mutation with genomic DNA from a plant homozygous for the wild-type 2490 gene. The probe for the Southern blot is prepared from a PCR product generated with the SLP369 (5' CAGACCACAATACCTTCAAAAATA 3': **SEQ ID NO:22**) and SLP370 (5' CCATTGTGTCTCCCTCCCGCTGTT 3': **SEQ ID NO:23**) primers. Finding an additional *Bam*H1 fragment in the 2490 heterozygote confirms a valid rescue.

The sequences obtained from the above clones are used in a BLASTn search against nucleotide sequence databases (Altschul et al. (1990) J Mol. Biol. 215: 403-410; Altschul et al (1997) Nucleic Acids Res. 25: 3389-3402). The search results show that the recovered sequences are identical to genomic DNA from *Arabidopsis* chromosome 5 P1 clone MTG13 (Genbank # AB008270). When the region of genomic DNA where the insertion event occurred is used in a BLASTn search of the Genbank EST database, four sequences derived from the ends of two ESTs, 144K24 (144K24 T7 Genbank #T76608 and 144K24XP Genbank #AA404903) and GBGF153 (5' end Genbank #F15182 and 3' end Genbank #F15181) are identified. The complete sequence of the 144K24 EST is determined and this sequence encodes the full open reading frame (ORF) for the 2490 gene. BLAST analysis of this EST indicates that the 2490 protein has sequence similarity with the *Brassica napus* Toc36 protein (Genbank #X79091; Ko et al. (1995) The Journal of Biological Chem. 270: 28601-28608; Wu et al. (1994) The Journal of Biological Chem. 269: 32264-32271; Pang et al. (1997) The Journal of Biological Chem. 272: 25623-25627). The Toc36 protein has also been referred to as bce44B, Com44, and Cim44. Because the genomic DNA that contains the 2490 ORF was not annotated correctly until now, the inventors are the first to provide experimental documentation of the correct ORF and sequence similarity for the 2490 gene.

**Example 4: Sequence Analysis of Tagged Seedling – Lethal Line #3963 From the T-DNA Mutagenized Population of *Arabidopsis***

The plasmid rescue technique is used to molecularly clone *Arabidopsis* genomic DNA flanking one or both sides of T-DNA insertions resulting from T-DNA mutagenesis. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of non-T-DNA vector sequences. The plasmids recovered from the plasmid rescue protocol are sequenced using the -21 primer (5' TGTAACGACGGCCAGT 3'; **SEQ ID NO:25**). Primer -21 provides information on the flanking sequence immediately adjacent to the right T-DNA border. Plasmid rescue is validated by PCR of genomic DNA from a heterozygote for the 3963 mutation. This PCR experiment uses a primer anchored in the predicted flanking sequence and the -21 primer (anchored in the T-DNA insertion). Finding a PCR product of the size expected based on the sequence of the plasmid rescued clone confirms a valid rescue. The sequence obtained from primer -21 is used in a BLASTn search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402.). The BLAST search results show that the recovered plant flanking sequence is 100% identical to the genomic sequence for P1 clone MDK4 on chromosome 5 (Genbank accession number AB010695). The T-DNA insertion occurred at base # 36342 of the annotated P1 clone MDK4 sequence, in the gene identified as MDK4.6. A tBLASTX analysis of the recovered flanking sequence shows sequence similarity to Mre11p, a DNA repair protein from *Saccharomyces cerevisiae* (Genbank accession number U60829). A fragment that encodes part of the *Arabidopsis* 3963 protein is isolated by amplification of *Arabidopsis* genomic DNA using the polymerase chain reaction. This fragment is used to probe an *Arabidopsis* cDNA library in the λYES vector (Elledge *et al.* (1991) Proc. Natl. Acad. Sci. 88:1731-1735). Positive phage clones are isolated and characterized using standard molecular biology techniques. The resultant cDNA clones are excised from the phage and the nucleotide sequence is determined. One cDNA clone is identified. The cDNA sequence is shown in **SEQ ID NO:7**. The deduced amino acid sequence is analyzed using the BLASTx search against nucleotide sequence databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25: 3389-3402). The BLAST search results show that the recovered 3963 cDNA shows sequence similarity to a number of DNA repair proteins, including Rad32p from *Schizosaccharomyces pombe* (Genbank accession number Q09683); hMre11 from *Homo sapiens* (Genbank accession number U37359); and Mre11p from *Saccharomyces*

*cerevisiae* (Genbank accession number U60829). Because the genomic DNA that contains the 3963 Open Reading Frame (ORF) was not annotated correctly in the prior art with respect to the exon/intron boundaries, the inventors are the first to provide experimental documentation of the correct ORF for the 3963 gene. The prior art indicates these exon/intron boundaries: 35662-35817, 36015-36172, 36315-36405, 36528-36647, 36728-36796, 36865-36956, 37045-37147, 37247-37354, 37476-37538, 37785-37862, 38060-38122, 38211-38271, 38753-38835, 38979-39092, 39468-39766, 39879-40002, 40161-40370. The exon/intron boundaries corresponding to the partial cDNA disclosed herein are: missing 5' end (first known base at 36147), 36147-36172, 36315-36405, 36528-36647, 36728-36796, 36865-36956, 37045-37147, 37247-37354, 37476-37538, 37610-37681, 37785-39092, 39212-39290, 39377-39445, 39532-39776, 39879-40002, 40161-40363, 40478-40508 (stop begins at 40509).

**Example 5: Sequence Analysis of Tagged Seedling – Lethal Line #4036 From the T-DNA Mutagenized Population of *Arabidopsis***

The plasmid rescue technique is used to molecularly clone *Arabidopsis* flanking DNA from one or both sides of the T-DNA insertions resulting from T-DNA mutagenesis. Plasmids obtained in this manner are analyzed by restriction enzyme digestion to sort the plasmids into classes based on their digestion pattern. For each class of plasmid clone, the DNA sequence is determined. The resulting sequences are analyzed for the presence of non-T-DNA vector sequences. The plasmids recovered from the plasmid rescue protocol are sequenced using the slp346 primer (5' GCGGACATCTACATTTTGA 3'; **SEQ ID NO:11**). Primer slp346 provides information on the flanking sequence immediately adjacent to the left T-DNA border. The plasmid rescue is validated via PCR of template genomic DNA from a heterozygote for the 4036 insertion mutation. The experiment uses a primer anchored in the predicted flanking sequence and the slp328 primer (5' ACCTTAGGCGACTTTTGAAC 3'; **SEQ ID NO:15**; anchored in the T-DNA insertion). Finding a PCR product of the size expected based on the sequence of the plasmid rescue clone confirms a valid rescue.

The sequence obtained from the above clone is used in a BLASTn search against nucleotide databases (Altschul *et al.* (1990) J Mol. Biol. 215:403-410; Altschul *et al.* (1997) Nucleic Acids Res. 25:3389-3402). The BLAST results show that the plant flanking sequence is 100% identical to published genomic sequence of P1 MQB2, from

chromosome 5 of *Arabidopsis* (Genbank accession # AB009053). The T-DNA insertion occurred at base 31,380 of the annotated P1 clone and interrupts a gene identified as MQB2.6. The protein encoded by the interrupted open reading frame (ORF) shows similarity to 1-deoxy-D-xylulose 5-phosphate reductoisomerase from a number of organisms including *Synechocystis* sp. (SWISS-PROT Q55663), *Bacillus subtilis* (SWISS-PROT O31753), and *Escherichia coli* (SWISS-PROT P45568) (Takahashi et al. (1998) Proc. Natl. Acad. Sci. USA, 95: 9879-9884). The genomic region encompassing the ORF is re-annotated with Web GeneMark software (Borodovsky, M. and McIninch J. (1993) Computers & Chemistry, 17: 123-133). Primers are then designed to the 5' and 3' ends of the predicted ORF, and PCR is performed using DNA from the pFL61 *Arabidopsis* cDNA library (Minet et al. (1992) Plant J. 2: 417-422) as the template. The resulting PCR product is TA-ligated and cloned (Original TA Cloning Kit, Invitrogen), and sequenced. Because the genomic DNA that contains the 4036 ORF was not annotated correctly in the prior art with respect to the exon/intron boundaries, the inventors are the first to provide experimental documentation of the correct ORF for the 4036 gene. The prior art indicates these exon/intron boundaries: 33490..33356, 31293..31207, 30971..30846, 30780..30718, 30622..30473, 30345..30288, 30194..30083, 29996..29892, 29805..29684, 29394..29248, 29162..28997. In the sequence of the present invention, base 31928 marks the first base of the cDNA's start codon and base 28996 marks the first base of the cDNA's stop codon. The 3' end of the exon containing the start codon is 31836, and the 5' end of the exon containing the stop codon is 29161. The internal exon/intron boundaries for the cDNA disclosed herein are: 31640.. 31448, 31294..31202, 30965..30843, 30777..30722, 30636..30473, 30355..30287, 30193..30082, 29995..29891, 29804..29684, 29394..29247.

#### Example 6a      Expression of Recombinant 245 Protein in *E. coli*

The coding region of the protein, corresponding to the cDNA clone SEQ ID NO: 1, is subcloned into previously described expression vectors, and transformed into *E. coli* using the manufacturer's conditions. Specific examples include plasmids such as pBluescript (Stratagene, La Jolla, CA), pFLAG (International Biotechnologies, Inc., New Haven, CT), and pTrcHis (Invitrogen, La Jolla, CA). *E. coli* is cultured, and expression of the 245 activity is confirmed. Protein conferring 245 activity is isolated using standard techniques.

**Example 6b      Expression of Recombinant 5283 Protein in *E. coli***

The coding region of the protein, corresponding to the cDNA clone SEQ ID NO: 3, is subcloned into previously described expression vectors, and transformed into *E. coli* using the manufacturer's conditions. Specific examples include plasmids such as pBluescript (Stratagene, La Jolla, CA), pFLAG (International Biotechnologies, Inc., New Haven, CT), and pTrcHis (Invitrogen, La Jolla, CA). *E. coli* is cultured, and expression of the 5283 activity is confirmed. Protein conferring 5283 activity is isolated using standard techniques.

**Example 6c      Expression of Recombinant 2490 Protein in *E. coli***

The coding region of the protein, corresponding to the cDNA clone SEQ ID NO: 5, is subcloned into previously described expression vectors, and transformed into *E. coli* using the manufacturer's conditions. Specific examples include plasmids such as pBluescript (Stratagene, La Jolla, CA), pFLAG (International Biotechnologies, Inc., New Haven, CT), and pTrcHis (Invitrogen, La Jolla, CA). *E. coli* is cultured, and expression of the 2490 activity is confirmed. Protein conferring 2490 activity is isolated using standard techniques.

**Example 6d      Expression of Recombinant 3963 Protein in *E. coli***

The coding region of the protein, corresponding to the cDNA clone SEQ ID NO: 7, is subcloned into previously described expression vectors, and transformed into *E. coli* using the manufacturer's conditions. Specific examples include plasmids such as pBluescript (Stratagene, La Jolla, CA), pFLAG (International Biotechnologies, Inc., New Haven, CT), and pTrcHis (Invitrogen, La Jolla, CA). *E. coli* is cultured, and expression of the 3963 activity is confirmed. Protein conferring 3963 activity is isolated using standard techniques.

**Example 6e      Expression of Recombinant 4036 Protein in *E. coli***

The coding region of the protein, corresponding to the cDNA clone SEQ ID NO: 9, is subcloned into previously described expression vectors, and transformed into *E. coli* using the manufacturer's conditions. Specific examples include plasmids such as pBluescript (Stratagene, La Jolla, CA), pFLAG (International Biotechnologies, Inc., New Haven, CT),

and pTrcHis (Invitrogen, La Jolla, CA). *E. coli* is cultured, and expression of the 4036 activity is confirmed. Protein conferring 4036 activity is isolated using standard techniques.

**Example 7: In vitro Recombination of 245, 5283, 2490, 3963, or 4036 Genes by DNA Shuffling**

The nucleotide sequence of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, respectively, is amplified by PCR. The resulting DNA fragment is digested by DNaseI treatment essentially as described (Stemmer et al. (1994) PNAS 91: 10747-10751) and the PCR primers are removed from the reaction mixture. A PCR reaction is carried out without primers and is followed by a PCR reaction with the primers, both as described (Stemmer et al. (1994) PNAS 91: 10747-10751). The resulting DNA fragments are cloned into pTRC99a (Pharmacia, Cat no: 27-5007-01) for use in bacteria, or into pESC vectors (Stratagene Catalog) for use in yeast; and transformed into a bacterial or yeast strain deficient in 245, 5283, 2490, 3963, or 4036 activity, respectively, by electroporation using the Biorad Gene Pulser and the manufacturer's conditions. The transformed bacteria or yeast are grown on medium that contains inhibitory concentrations of an inhibitor of 245, 5283, 2490, 3963, or 4036 activity and those colonies that grow in the presence of the inhibitor are selected. Colonies that grow in the presence of normally inhibitory concentrations of inhibitor are picked and purified by repeated restreaking. Their plasmids are purified and the DNA sequences of cDNA inserts from plasmids that pass this test are then determined.

In a similar reaction, PCR-amplified DNA fragments comprising the *A. thaliana* 245, 5283, 2490, 3963, or 4036 gene, respectively, encoding the protein and PCR-amplified DNA fragments comprising the 245, 5283, 2490, 3963, or 4036 gene, respectively, from *E. coli* are recombined *in vitro* and resulting variants with improved tolerance to the inhibitor are recovered as described above.

**Example 8a: In vitro Recombination of 245 Genes by Staggered Extension Process**

The *A. thaliana* 245 gene encoding the 245 protein and the *E.coli* 245 gene are each cloned into the polylinker of a pBluescript vector. A PCR reaction is carried out essentially as described (Zhao et al. (1998) Nature Biotechnology 16: 258-261) using the "reverse primer" and the "M13 -20 primer" (Stratagene Catalog). Amplified PCR fragments are digested with appropriate restriction enzymes and cloned into pTRC99a and mutated 245 genes are screened as described in Example 7.

**Example 8b: In vitro Recombination of 5283 Genes by Staggered Extension Process**

The *A. thaliana* 5283 gene encoding the 5283 protein and the *E.coli* 5283 gene are each cloned into the polylinker of a pBluescript vector. A PCR reaction is carried out essentially as described (Zhao et al. (1998) Nature Biotechnology 16: 258-261) using the "reverse primer" and the "M13 -20 primer" (Stratagene Catalog). Amplified PCR fragments are digested with appropriate restriction enzymes and cloned into pTRC99a and mutated 5283 genes are screened as described in Example 7.

**Example 8c: In vitro Recombination of 2490 Genes by Staggered Extension Process**

The *A. thaliana* 2490 gene encoding the 2490 protein and the *E.coli* 2490 gene are each cloned into the polylinker of a pBluescript vector. A PCR reaction is carried out essentially as described (Zhao et al. (1998) Nature Biotechnology 16: 258-261) using the "reverse primer" and the "M13 -20 primer" (Stratagene Catalog). Amplified PCR fragments are digested with appropriate restriction enzymes and cloned into pTRC99a and mutated 2490 genes are screened as described in Example 7.

**Example 8d: In vitro Recombination of 3963 Genes by Staggered Extension Process**

The *A. thaliana* 3963 gene encoding the 3963 protein and the *E.coli* 3963 gene are each cloned into the polylinker of a pBluescript vector. A PCR reaction is carried out essentially as described (Zhao et al. (1998) Nature Biotechnology 16: 258-261) using the "reverse primer" and the "M13 -20 primer" (Stratagene Catalog). Amplified PCR fragments are digested with appropriate restriction enzymes and cloned into pTRC99a and mutated 3963 genes are screened as described in Example 7.



**Example 8e: In vitro Recombination of 4036 Genes by Staggered Extension Process**

The *A. thaliana* 4036 gene encoding the 4036 protein and the *E.coli* 4036 gene are each cloned into the polylinker of a pBluescript vector. A PCR reaction is carried out essentially as described (Zhao et al. (1998) Nature Biotechnology 16: 258-261) using the "reverse primer" and the "M13 -20 primer" (Stratagene Catalog). Amplified PCR fragments are digested with appropriate restriction enzymes and cloned into pTRC99a and mutated 4036 genes are screened as described in Example 7.

**Example 9: In Vitro Binding Assays**

Recombinant 245, 5283, 2490, 3963, or 4036 protein is obtained, for example, according to Example 6a,6b,6c,6d,or 6e, respectively. The protein is immobilized on chips appropriate for ligand binding assays using techniques which are well known in the art. The protein immobilized on the chip is exposed to sample compound in solution according to methods well know in the art. While the sample compound is in contact with the immobilized protein measurements capable of detecting protein-ligand interactions are conducted. Examples of such measurements are SELDI, biacore and FCS, described above. Compounds found to bind the protein are readily discovered in this fashion and are subjected to further characterization.

The above disclosed embodiments are illustrative. This disclosure of the invention will place one skilled in the art in possession of many variations of the invention. All such obvious and foreseeable variations are intended to be encompassed by the appended claims.

**What Is Claimed Is:**

1. An isolated DNA molecule comprising a nucleotide sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.
2. The DNA molecule of claim 1, wherein the sequence encodes an amino acid sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10.
3. The DNA molecule of claim 1, wherein the sequence is any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.
4. The DNA molecule of claim 1, wherein the sequence encodes the amino acid sequence of any one of the sequences selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10.
5. The DNA molecule according to claim 1, wherein said nucleotide sequence is a plant nucleotide sequence.
6. The DNA molecule of claim 5, wherein the plant is *Arabidopsis thaliana*.
7. The DNA molecule of claim 1, wherein the protein has any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity.
8. An amino acid sequence comprising an amino acid sequence encoded by a nucleotide sequence substantially similar to any one of the sequences selected from the group

consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.

9. The amino acid sequence of claim 8 comprising an amino acid sequence encoded by any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.
10. An amino acid sequence comprising an amino acid sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10.
11. The amino acid sequence of claim 10, wherein the sequence is any one of the sequences selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10.
12. The amino acid sequence of claim 8, wherein the protein has any one of the activities selected from the group of 245, 5283, 2490, 3963 and 4036 activity.
13. An amino acid sequence comprising at least 20 consecutive amino acid residues of the amino acid sequence encoded by any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.
14. An amino acid sequence comprising at least 20 consecutive amino acid residues of the amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8 and SEQ ID NO:10.
15. An expression cassette comprising a promoter operatively linked to a DNA molecule according to claim 1.
16. A recombinant vector comprising an expression cassette according to claim 15, wherein said vector is capable of being stably transformed into a host cell.

17. A host cell comprising an expression cassette according to claim 15, wherein said nucleotide sequence is expressible in said cell.
18. A host cell according to claim 17, wherein said host cell is an eukaryotic cell.
19. A host cell according to claim 17, wherein said host cell is selected from the group consisting of an insect cell, a yeast cell, and a plant cell.
20. A host cell according to claim 17, wherein said host cell is a prokaryotic cell.
21. A host cell according to claim 17, wherein said host cell is a bacterial cell.
22. A plant or seed comprising a plant cell of claim 19.
23. A plant of claim 22, wherein said plant is tolerant to an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity.
24. A process for making nucleotides sequences encoding gene products having altered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity comprising,
  - a) shuffling a nucleotide sequence of claim 1,
  - b) expressing the resulting shuffled nucleotide sequences and
  - c) selecting for altered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity as compared to the activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity of the gene product of said unmodified nucleotide sequence.
25. The process of claim 24, wherein the nucleotide sequence is any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9.

26. A shuffled DNA molecule obtainable by the process of claim 24.
27. A shuffled DNA molecule produced by the process of claim 24.
28. A shuffled DNA molecule obtained by the process of claim 24, wherein said shuffled DNA molecule encodes a gene product having enhanced tolerance to an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity.
29. An expression cassette comprising a promoter operatively linked to a nucleotide sequence according to claim 26.
30. A recombinant vector comprising an expression cassette according to claim 29, wherein said vector is capable of being stably transformed into a host cell.
31. A host cell comprising an expression cassette according to claim 29, wherein said nucleotide sequence is expressible in said cell.
32. A host cell according to claim 31, wherein said host cell is an eukaryotic cell.
33. A host cell according to claim 31, wherein said host cell is selected from the group consisting of an insect cell, a yeast cell, and a plant cell.
34. A host cell according to claim 31, wherein said host cell is a prokaryotic cell.
35. A host cell according to claim 31, wherein said host cell is a bacterial cell.
36. A plant or seed comprising a plant cell of claim 33.
37. A plant of claim 36, wherein said plant is tolerant to an inhibitor selected from the group consisting of 245, 5283, 2490, 3963 and 4036 activity.
38. A method for selecting compounds that interact with the protein encoded by any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ

ID NO:5, SEQ ID NO:7 and SEQ ID NO:9. ????

, comprising:

a) expressing a DNA molecule comprising any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9 or a sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9 to generate the corresponding protein,

b) testing a compound suspected of having the ability to interact with the protein expressed in step (a), and

c) selecting compounds that interact with the protein in step (b).

39. A process of identifying an inhibitor of any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity comprising:

a) introducing a DNA molecule comprising a nucleotide sequence of any one of the sequences selected from the group consisting of SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7 and SEQ ID NO:9 and having any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity, or nucleotide sequences substantially similar thereto, or a homolog thereof, into a plant cell, such that said sequence is functionally expressible at levels that are higher than wild-type expression levels,

b) combining said plant cell with a compound to be tested for the ability to inhibit any one of the activities selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity under conditions conducive to such inhibition,

- c) measuring plant cell growth under the conditions of step (b), and
  - d) comparing the growth of said plant cell with the growth of a plant cell having an unaltered activity selected from the group consisting of 245 activity, 5283 activity, 2490 activity, 3963 activity and 4036 activity under identical conditions, and
  - e) selecting said compound that inhibits plant cell growth in step (d).
40. A compound having herbicidal activity identifiable according to the process of claim 39.
41. A process of identifying compounds having herbicidal activity comprising:
- a) combining a protein of claim 8 and a compound to be tested for the ability to interact with said protein, under conditions conducive to interaction,
  - b) selecting a compound identified in step (a) that is capable of interacting with said protein,
  - c) applying identified compound in step (b) to a plant to test for herbicidal activity, and
  - d) selecting compounds having herbicidal activity.
42. A compound having herbicidal activity identifiable according to the process of claim 41.
43. A method for suppressing the growth of a plant comprising, applying to said plant a compound that inhibits the activity of the amino acid sequence of claim 8 in an amount sufficient to suppress the growth of said plant.
44. The method of claim 41, wherein the compound is a compound having herbicidal activity identifiable according to the process of claim 39.

45. A method of improving crops comprising, applying to a herbicide tolerant plant or seed selected from the group consisting of the plant or seed of claim 23 and the plant or seed of claim 37, a compound having herbicidal activity identifiable according to a process selected from the group of the method of claim 38, the process of claim 39, and the process of claim 41, in an amount that inhibits the growth of undesired vegetation without significantly suppressing the growth of the herbicide tolerant plant or seed.

46. A DNA molecule comprising a nucleotide sequence substantially similar to any one of the sequences selected from the group consisting of SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29.



## SEQUENCE LISTING

&lt;110&gt; Novartis AG

&lt;120&gt; Herbicide Target Genes and Methods

&lt;130&gt; combined herbicide target genes

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 29

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1119

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1119)

&lt;400&gt; 1

atg gat gac atg gac acc gtc tac aag caa ttg gga ttg ttt tca cta	48
Met Asp Asp Met Asp Thr Val Tyr Lys Gln Leu Gly Leu Phe Ser Leu	
1 5 10 15	
aag aag aag att aaa gat gtt gtt ctt aag gct gag atg ttt gca ccg	96
Lys Lys Lys Ile Lys Asp Val Val Leu Lys Ala Glu Met Phe Ala Pro	
20 25 30	
gat gct ctt gag ctt gaa gaa gag cag tgg ata aag caa gaa gaa aca	144
Asp Ala Leu Glu Leu Glu Glu Glu Gln Trp Ile Lys Gln Glu Glu Thr	
35 40 45	
atg cgt tac ttt gat tta tgg gat gat ccc gct aaa tct gat gag att	192
Met Arg Tyr Phe Asp Leu Trp Asp Asp Pro Ala Lys Ser Asp Glu Ile	
50 55 60	
ctt ctc aaa tta gct gat cga gct aaa gca gtc gat tcc ctc aaa gac	240
Leu Leu Lys Leu Ala Asp Arg Ala Lys Ala Val Asp Ser Leu Lys Asp	
65 70 75 80	
ctc aaa tac aag gct gaa gaa gct aag ctg atc ata caa ttg ggt gag	288
Leu Lys Tyr Lys Ala Glu Glu Ala Lys Leu Ile Ile Gln Leu Gly Glu	
85 90 95	
atg gat gct ata gat tac agt ctc ttt gag caa gcc tat gat tca tca	336
Met Asp Ala Ile Asp Tyr Ser Leu Phe Glu Gln Ala Tyr Asp Ser Ser	
100 105 110	
ctc gat gta agt aga tcg ttg cat cac tat gag atg tct aag ctt ctt	384
Leu Asp Val Ser Arg Ser Leu His His Tyr Glu Met Ser Lys Leu Leu	

115	120	125	
agg gat caa tat gac gct gaa ggc gct tgt atg att atc aaa tct gga			432
Arg Asp Gln Tyr Asp Ala Glu Gly Ala Cys Met Ile Ile Lys Ser Gly			
130	135	140	
tct cca ggc gca aaa tct cag ata tgg aca gag caa gtt gta agt atg			480
Ser Pro Gly Ala Lys Ser Gln Ile Trp Thr Glu Gln Val Val Ser Met			
145	150	155	160
tat atc aaa tgg gca gaa agg cta ggc caa aac gcg cgg gtg gct gag			528
Tyr Ile Lys Trp Ala Glu Arg Leu Gly Gln Asn Ala Arg Val Ala Glu			
165	170	175	
aaa tgt agt tta ttg agt aat aaa agt ggc gta agt tca gcc acg ata			576
Lys Cys Ser Leu Leu Ser Asn Lys Ser Gly Val Ser Ser Ala Thr Ile			
180	185	190	
gag ttt gaa ttc gag ttt gct tat ggt tat ctc tta ggt gag cga ggt			624
Glu Phe Glu Phe Glu Phe Ala Tyr Gly Tyr Leu Leu Gly Glu Arg Gly			
195	200	205	
gtg cac cgc ctt atc ata agt tcc act tct aat gag gaa tgt tca gcg			672
Val His Arg Leu Ile Ile Ser Ser Thr Ser Asn Glu Glu Cys Ser Ala			
210	215	220	
act gtt gat atc ata cca cta ttc ttg aga gca tct cct gat ttt gaa			720
Thr Val Asp Ile Ile Pro Leu Phe Leu Arg Ala Ser Pro Asp Phe Glu			
225	230	235	240
gta aag gaa ggt gat ttg att gta tcg tat cct gca aaa gag gat cac			768
Val Lys Glu Gly Asp Leu Ile Val Ser Tyr Pro Ala Lys Glu Asp His			
245	250	255	
aaa ata gct gag aat atg gtt tgt atc cac cat att ccg agt gga gta			816
Lys Ile Ala Glu Asn Met Val Cys Ile His His Ile Pro Ser Gly Val			
260	265	270	
aca cta caa tct tca gga gaa aga aac cgg ttt gca aac agg atc aaa			864
Thr Leu Gln Ser Ser Gly Glu Arg Asn Arg Phe Ala Asn Arg Ile Lys			
275	280	285	
gct cta aac cgg ttg aag gcg aag cta ctt gtg ata gca aaa gag caa			912
Ala Leu Asn Arg Leu Lys Ala Lys Leu Leu Val Ile Ala Lys Glu Gln			
290	295	300	
aag gtt tcg gat gta aat aaa atc gac agc aag aac att ttg gaa ccg			960
Lys Val Ser Asp Val Asn Lys Ile Asp Ser Lys Asn Ile Leu Glu Pro			
305	310	315	320
cgg gaa gaa acc agg agt tat gtc tct aag ggt cac aag atg gtg gtt			1008
Arg Glu Glu Thr Arg Ser Tyr Val Ser Lys Gly His Lys Met Val Val			
325	330	335	
gat aga aaa acc ggt tta gag att ctg gac ctg aaa tcg gtc ttg gat			1056

Asp Arg Lys Thr Gly Leu Glu Ile Leu Asp Leu Lys Ser Val Leu Asp  
 340 345 350

gga aac att gga cca ctc ctt gga gct cat att agc atg aga aga tca 1104  
 Gly Asn Ile Gly Pro Leu Leu Gly Ala His Ile Ser Met Arg Arg Ser  
 355 360 365

att gat gcg att tag 1119  
 Ile Asp Ala Ile  
 370

<210> 2

<211> 372

<212> PRT

<213> Arabidopsis thaliana

<400> 2

Met Asp Asp Met Asp Thr Val Tyr Lys Gln Leu Gly Leu Phe Ser Leu  
 1 5 10 15  
 Lys Lys Lys Ile Lys Asp Val Val Leu Lys Ala Glu Met Phe Ala Pro  
 20 25 30  
 Asp Ala Leu Glu Leu Glu Glu Glu Trp Ile Lys Gln Glu Glu Thr  
 35 40 45  
 Met Arg Tyr Phe Asp Leu Trp Asp Asp Pro Ala Lys Ser Asp Glu Ile  
 50 55 60  
 Leu Leu Lys Leu Ala Asp Arg Ala Lys Ala Val Asp Ser Leu Lys Asp  
 65 70 75 80  
 Leu Lys Tyr Lys Ala Glu Glu Ala Lys Leu Ile Ile Gln Leu Gly Glu  
 85 90 95  
 Met Asp Ala Ile Asp Tyr Ser Leu Phe Glu Gln Ala Tyr Asp Ser Ser  
 100 105 110  
 Leu Asp Val Ser Arg Ser Leu His His Tyr Glu Met Ser Lys Leu Leu  
 115 120 125  
 Arg Asp Gln Tyr Asp Ala Glu Gly Ala Cys Met Ile Ile Lys Ser Gly  
 130 135 140  
 Ser Pro Gly Ala Lys Ser Gln Ile Trp Thr Glu Gln Val Val Ser Met  
 145 150 155 160  
 Tyr Ile Lys Trp Ala Glu Arg Leu Gly Gln Asn Ala Arg Val Ala Glu  
 165 170 175  
 Lys Cys Ser Leu Leu Ser Asn Lys Ser Gly Val Ser Ser Ala Thr Ile  
 180 185 190  
 Glu Phe Glu Phe Glu Phe Ala Tyr Gly Tyr Leu Leu Gly Glu Arg Gly  
 195 200 205  
 Val His Arg Leu Ile Ile Ser Ser Thr Ser Asn Glu Glu Cys Ser Ala  
 210 215 220  
 Thr Val Asp Ile Ile Pro Leu Phe Leu Arg Ala Ser Pro Asp Phe Glu  
 225 230 235 240  
 Val Lys Glu Gly Asp Leu Ile Val Ser Tyr Pro Ala Lys Glu Asp His  
 245 250 255  
 Lys Ile Ala Glu Asn Met Val Cys Ile His His Ile Pro Ser Gly Val  
 260 265 270  
 Thr Leu Gln Ser Ser Gly Glu Arg Asn Arg Phe Ala Asn Arg Ile Lys  
 275 280 285  
 Ala Leu Asn Arg Leu Lys Ala Lys Leu Leu Val Ile Ala Lys Glu Gln

```

      290              295              300
Lys Val Ser Asp Val Asn Lys Ile Asp Ser Lys Asn Ile Leu Glu Pro
305              310              315              320
Arg Glu Glu Thr Arg Ser Tyr Val Ser Lys Gly His Lys Met Val Val
      325              330              335
Asp Arg Lys Thr Gly Leu Glu Ile Leu Asp Leu Lys Ser Val Leu Asp
      340              345              350
Gly Asn Ile Gly Pro Leu Leu Gly Ala His Ile Ser Met Arg Arg Ser
      355              360              365
Ile Asp Ala Ile
      370

```

<210> 3  
 <211> 1458  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (1)..(1458)

```

<400> 3
atg gca act ctt gaa gat tct ttc ctt gct gat ttg gac gag tta tct 48
Met Ala Thr Leu Glu Asp Ser Phe Leu Ala Asp Leu Asp Glu Leu Ser
      1              5              10              15

gac aat gaa gca gaa ttg gac gag aat gat ggt gat gtt gga aag gaa 96
Asp Asn Glu Ala Glu Leu Asp Glu Asn Asp Gly Asp Val Gly Lys Glu
      20              25              30

gaa gaa gat gtt gat atg gat atg gct gat tta gag aca ctt aac tat 144
Glu Glu Asp Val Asp Met Asp Met Ala Asp Leu Glu Thr Leu Asn Tyr
      35              40              45

gat gat ctc gat aat gtt tct aag ctg cag aag agt cag aga tat gct 192
Asp Asp Leu Asp Asn Val Ser Lys Leu Gln Lys Ser Gln Arg Tyr Ala
      50              55              60

gat att atg cat aaa gta gag gag gct ctt ggg aaa gat tct gat gga 240
Asp Ile Met His Lys Val Glu Glu Ala Leu Gly Lys Asp Ser Asp Gly
      65              70              75              80

gct gag aaa gga act gtc ttg gaa gat gat cct gag tat aag ctt att 288
Ala Glu Lys Gly Thr Val Leu Glu Asp Asp Pro Glu Tyr Lys Leu Ile
      85              90              95

gtg gat tgt aat cag ctt tcg gtc gat att gag aat gaa atc gtt att 336
Val Asp Cys Asn Gln Leu Ser Val Asp Ile Glu Asn Glu Ile Val Ile
      100             105             110

gtc cac aac ttt atc aaa gac aag tac aag ctt aag ttt caa gag ctt 384
Val His Asn Phe Ile Lys Asp Lys Tyr Lys Leu Lys Phe Gln Glu Leu
      115             120             125

```

gag tcg ttg gtt cat cac cct att gac tat gca tgt gtt gtg aag aag Glu Ser Leu Val His His Pro Ile Asp Tyr Ala Cys Val Val Lys Lys 130 135 140	432
att ggg aat gag acg gat ttg gct ctt gtt gat ctc gct gac ctt ctt Ile Gly Asn Glu Thr Asp Leu Ala Leu Val Asp Leu Ala Asp Leu Leu 145 150 155 160	480
cct tca gct att atc atg gtt gtt tca gtt act gct tta act acg aaa Pro Ser Ala Ile Ile Met Val Val Ser Val Thr Ala Leu Thr Thr Lys 165 170 175	528
ggg agt gca ctg cca gag gat gtt ttg caa aag gtg tta gag gct tgt Gly Ser Ala Leu Pro Glu Asp Val Leu Gln Lys Val Leu Glu Ala Cys 180 185 190	576
gat cgg gct tta gat ctt gat tcc gca agg aag aag gtc ctt gag ttt Asp Arg Ala Leu Asp Leu Asp Ser Ala Arg Lys Lys Val Leu Glu Phe 195 200 205	624
gtt gaa agt aag atg gga tct att gca cct aat ctt tct gct att gtt Val Glu Ser Lys Met Gly Ser Ile Ala Pro Asn Leu Ser Ala Ile Val 210 215 220	672
ggg agt gct gtt gca gcc aaa ctc atg ggg act gct gga ggt ttg tca Gly Ser Ala Val Ala Ala Lys Leu Met Gly Thr Ala Gly Gly Leu Ser 225 230 235 240	720
gca ctt gct aaa atg cct gcg tgt aat gtt caa gtt ctt ggc cac aag Ala Leu Ala Lys Met Pro Ala Cys Asn Val Gln Val Leu Gly His Lys 245 250 255	768
agg aag aac ctt gct ggg ttt tct tct gca acg tct cag tcc cgt gtg Arg Lys Asn Leu Ala Gly Phe Ser Ser Ala Thr Ser Gln Ser Arg Val 260 265 270	816
ggt tat ctg gag cag aca gag att tac caa agc acg cct cct gga ctt Gly Tyr Leu Glu Gln Thr Glu Ile Tyr Gln Ser Thr Pro Pro Gly Leu 275 280 285	864
cag gct cgc gct ggc agg ctc gtg gct gca aaa tca act ttg gca gca Gln Ala Arg Ala Gly Arg Leu Val Ala Ala Lys Ser Thr Leu Ala Ala 290 295 300	912
aga gtt gat gct act aga ggg gat ccg tta ggg ata agt gga aaa gct Arg Val Asp Ala Thr Arg Gly Asp Pro Leu Gly Ile Ser Gly Lys Ala 305 310 315 320	960
ttc agg gag gag atc cgt aag aag att gag aaa tgg caa gaa cct cct Phe Arg Glu Glu Ile Arg Lys Lys Ile Glu Lys Trp Gln Glu Pro Pro 325 330 335	1008
cct gca aga cag cct aag cca ctt cct gtt cct gat tct gaa ccg aag Pro Ala Arg Gln Pro Lys Pro Leu Pro Val Pro Asp Ser Glu Pro Lys	1056

340	345	350	
aaa aga agg ggt ggt cgc cgt cta aga aaa atg aaa gaa agg tat caa			1104
Lys Arg Arg Gly Gly Arg Arg Leu Arg Lys Met Lys Glu Arg Tyr Gln			
355	360	365	
gta aca gat atg agg aag ctg gcc aac aga atg gcg ttt ggt aca cct			1152
Val Thr Asp Met Arg Lys Leu Ala Asn Arg Met Ala Phe Gly Thr Pro			
370	375	380	
gaa gag agc tcc ctc ggt gat gga cta gga gaa ggt tat gga atg ctt			1200
Glu Glu Ser Ser Leu Gly Asp Gly Leu Gly Glu Gly Tyr Gly Met Leu			
385	390	395	400
ggc cag gca gga agc aac agg ctg cga gta tcc agt gtt ccg agc aag			1248
Gly Gln Ala Gly Ser Asn Arg Leu Arg Val Ser Ser Val Pro Ser Lys			
405	410	415	
ctt aag att aat gct aag gtc gcc aaa aag ctt aaa gaa agg cag tat			1296
Leu Lys Ile Asn Ala Lys Val Ala Lys Lys Leu Lys Glu Arg Gln Tyr			
420	425	430	
gcg ggt ggt gcg act acc tct ggt ttg aca tog agc ctg gct ttc act			1344
Ala Gly Gly Ala Thr Thr Ser Gly Leu Thr Ser Ser Leu Ala Phe Thr			
435	440	445	
cct gtg cag gga ata gag ttg tgc aat cct cag cag gct tta gga tta			1392
Pro Val Gln Gly Ile Glu Leu Cys Asn Pro Gln Gln Ala Leu Gly Leu			
450	455	460	
gga agt ggg act caa agc act tac ttc tca gag tca gga acc ttc tcg			1440
Gly Ser Gly Thr Gln Ser Thr Tyr Phe Ser Glu Ser Gly Thr Phe Ser			
465	470	475	480
aag ctg aag aag atc taa			1458
Lys Leu Lys Lys Ile			
485			

&lt;210&gt; 4

&lt;211&gt; 485

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 4

Met Ala Thr Leu Glu Asp Ser Phe Leu Ala Asp Leu Asp Glu Leu Ser			
1	5	10	15
Asp Asn Glu Ala Glu Leu Asp Glu Asn Asp Gly Asp Val Gly Lys Glu			
20	25	30	
Glu Glu Asp Val Asp Met Asp Met Ala Asp Leu Glu Thr Leu Asn Tyr			
35	40	45	
Asp Asp Leu Asp Asn Val Ser Lys Leu Gln Lys Ser Gln Arg Tyr Ala			
50	55	60	
Asp Ile Met His Lys Val Glu Glu Ala Leu Gly Lys Asp Ser Asp Gly			
65	70	75	80

Ala	Glu	Lys	Gly	Thr	Val	Leu	Glu	Asp	Pro	Glu	Tyr	Lys	Leu	Ile
				85					90				95	
Val	Asp	Cys	Asn	Gln	Leu	Ser	Val	Asp	Ile	Glu	Asn	Glu	Ile	Val
			100					105					110	Ile
Val	His	Asn	Phe	Ile	Lys	Asp	Lys	Tyr	Lys	Leu	Lys	Phe	Gln	Glu
		115				120						125		Leu
Glu	Ser	Leu	Val	His	His	Pro	Ile	Asp	Tyr	Ala	Cys	Val	Val	Lys
	130					135					140			Lys
Ile	Gly	Asn	Glu	Thr	Asp	Leu	Ala	Leu	Val	Asp	Leu	Ala	Asp	Leu
145					150					155				160
Pro	Ser	Ala	Ile	Ile	Met	Val	Val	Ser	Val	Thr	Ala	Leu	Thr	Thr
				165					170					175
Gly	Ser	Ala	Leu	Pro	Glu	Asp	Val	Leu	Gln	Lys	Val	Leu	Glu	Ala
			180					185					190	Cys
Asp	Arg	Ala	Leu	Asp	Leu	Asp	Ser	Ala	Arg	Lys	Lys	Val	Leu	Glu
	195					200						205		Phe
Val	Glu	Ser	Lys	Met	Gly	Ser	Ile	Ala	Pro	Asn	Leu	Ser	Ala	Ile
	210					215				220				Val
Gly	Ser	Ala	Val	Ala	Ala	Lys	Leu	Met	Gly	Thr	Ala	Gly	Gly	Leu
225					230					235				240
Ala	Leu	Ala	Lys	Met	Pro	Ala	Cys	Asn	Val	Gln	Val	Leu	Gly	His
				245				250						255
Arg	Lys	Asn	Leu	Ala	Gly	Phe	Ser	Ser	Ala	Thr	Ser	Gln	Ser	Arg
		260						265					270	Val
Gly	Tyr	Leu	Glu	Gln	Thr	Glu	Ile	Tyr	Gln	Ser	Thr	Pro	Pro	Gly
	275					280						285		Leu
Gln	Ala	Arg	Ala	Gly	Arg	Leu	Val	Ala	Ala	Lys	Ser	Thr	Leu	Ala
	290					295					300			
Arg	Val	Asp	Ala	Thr	Arg	Gly	Asp	Pro	Leu	Gly	Ile	Ser	Gly	Lys
305					310					315				320
Phe	Arg	Glu	Glu	Ile	Arg	Lys	Lys	Ile	Glu	Lys	Trp	Gln	Glu	Pro
				325				330						335
Pro	Ala	Arg	Gln	Pro	Lys	Pro	Leu	Pro	Val	Pro	Asp	Ser	Glu	Pro
			340					345					350	Lys
Lys	Arg	Arg	Gly	Gly	Arg	Arg	Leu	Arg	Lys	Met	Lys	Glu	Arg	Tyr
		355					360					365		Gln
Val	Thr	Asp	Met	Arg	Lys	Leu	Ala	Asn	Arg	Met	Ala	Phe	Gly	Thr
	370					375					380			Pro
Glu	Glu	Ser	Ser	Leu	Gly	Asp	Gly	Leu	Gly	Glu	Gly	Tyr	Gly	Met
385					390					395				400
Gly	Gln	Ala	Gly	Ser	Asn	Arg	Leu	Arg	Val	Ser	Ser	Val	Pro	Ser
				405					410					Lys
Leu	Lys	Ile	Asn	Ala	Lys	Val	Ala	Lys	Lys	Leu	Lys	Glu	Arg	Gln
			420					425					430	Tyr
Ala	Gly	Gly	Ala	Thr	Thr	Ser	Gly	Leu	Thr	Ser	Ser	Leu	Ala	Phe
		435					440					445		Thr
Pro	Val	Gln	Gly	Ile	Glu	Leu								

<210> 5  
 <211> 1344  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (1)..(1344)

<400> 5  
 atg gag aac ctt acc cta gtt tct tgc tca gct tct tct cca aag ctg 48  
 Met Glu Asn Leu Thr Leu Val Ser Cys Ser Ala Ser Ser Pro Lys Leu  
 1 5 10 15  
 tta att gga tgc aat ttc act tcc tcg ctg aaa aac cct act ggg ttt 96  
 Leu Ile Gly Cys Asn Phe Thr Ser Ser Leu Lys Asn Pro Thr Gly Phe  
 20 25 30  
 tct cgt cgg act cct aat att gtc ctc cgg tgt tcc aaa ata tct gcc 144  
 Ser Arg Arg Thr Pro Asn Ile Val Leu Arg Cys Ser Lys Ile Ser Ala  
 35 40 45  
 tct gct caa tct caa tct ccc tct tcg cgt ccg gag aac act gga gaa 192  
 Ser Ala Gln Ser Gln Ser Pro Ser Ser Arg Pro Glu Asn Thr Gly Glu  
 50 55 60  
 atc gtg gtt gtg aaa cag aga agc aaa gct ttt gca agt ata ttt tct 240  
 Ile Val Val Val Lys Gln Arg Ser Lys Ala Phe Ala Ser Ile Phe Ser  
 65 70 75 80  
 tcg agt cgt gat caa cag aca act tct gtt gct tcc cct agt gtg cct 288  
 Ser Ser Arg Asp Gln Gln Thr Thr Ser Val Ala Ser Pro Ser Val Pro  
 85 90 95  
 gtg cca cca cca tct tca tca acc ata gga tca cca ctt ttc tgg att 336  
 Val Pro Pro Pro Ser Ser Ser Thr Ile Gly Ser Pro Leu Phe Trp Ile  
 100 105 110  
 ggt gtt ggt gtt ggt cta tca gct ttg ttc tca tat gta act tca aat 384  
 Gly Val Gly Val Gly Leu Ser Ala Leu Phe Ser Tyr Val Thr Ser Asn  
 115 120 125  
 tta aag aaa tat gca atg caa aca gct atg aag acg atg atg aac caa 432  
 Leu Lys Lys Tyr Ala Met Gln Thr Ala Met Lys Thr Met Met Asn Gln  
 130 135 140  
 atg aat acg caa aat agc cag ttt aat aat tct gga ttc cca tca gga 480  
 Met Asn Thr Gln Asn Ser Gln Phe Asn Asn Ser Gly Phe Pro Ser Gly  
 145 150 155 160  
 tca cct ttt ccg ttt cca ttt cct cct caa aca agt cct gct tcc tcg 528  
 Ser Pro Phe Pro Phe Pro Phe Pro Pro Gln Thr Ser Pro Ala Ser Ser  
 165 170 175  
 cca ttc caa tct caa tcc cag tct tca ggt gct acc gtt gat gtg aca 576



Pro	Phe	Gln	Ser	Gln	Ser	Gln	Ser	Ser	Gly	Ala	Thr	Val	Asp	Val	Thr	
		180						185					190			
gcg	aca	aaa	gta	gag	aca	cct	cct	tca	act	aaa	ccg	aaa	cct	aca	cct	624
Ala	Thr	Lys	Val	Glu	Thr	Pro	Pro	Ser	Thr	Lys	Pro	Lys	Pro	Thr	Pro	
		195					200					205				
gca	aag	gat	ata	gag	gtg	gat	aag	cca	agt	gtt	gtc	tta	gag	gca	agc	672
Ala	Lys	Asp	Ile	Glu	Val	Asp	Lys	Pro	Ser	Val	Val	Leu	Glu	Ala	Ser	
	210					215					220					
aaa	gag	aag	aaa	gaa	gaa	aag	aac	tat	gcc	ttt	gaa	gac	att	tca	ccc	720
Lys	Glu	Lys	Lys	Glu	Glu	Lys	Asn	Tyr	Ala	Phe	Glu	Asp	Ile	Ser	Pro	
225				230					235					240		
gag	gaa	acc	aca	aaa	gaa	agc	cca	ttt	agc	aac	tat	gca	gaa	gtc	tct	768
Glu	Glu	Thr	Thr	Lys	Glu	Ser	Pro	Phe	Ser	Asn	Tyr	Ala	Glu	Val	Ser	
			245					250						255		
gaa	act	aat	tcc	ccc	aaa	gaa	act	cgc	ttg	ttt	gag	gat	gtc	ttg	caa	816
Glu	Thr	Asn	Ser	Pro	Lys	Glu	Thr	Arg	Leu	Phe	Glu	Asp	Val	Leu	Gln	
		260						265					270			
aat	gga	gct	ggc	ccg	gca	aat	ggc	gcc	act	gct	tca	gag	gtt	ttt	caa	864
Asn	Gly	Ala	Gly	Pro	Ala	Asn	Gly	Ala	Thr	Ala	Ser	Glu	Val	Phe	Gln	
	275					280						285				
tct	ttg	ggc	ggc	ggg	aaa	gga	ggg	ccg	ggc	ttc	tct	gta	gaa	gct	tta	912
Ser	Leu	Gly	Gly	Gly	Lys	Gly	Gly	Pro	Gly	Leu	Ser	Val	Glu	Ala	Leu	
	290				295					300						
gag	aaa	atg	atg	gaa	gat	cca	aca	gtc	cag	aag	atg	gtt	tac	cca	tac	960
Glu	Lys	Met	Met	Glu	Asp	Pro	Thr	Val	Gln	Lys	Met	Val	Tyr	Pro	Tyr	
305				310					315					320		
ttg	cct	gag	gag	atg	agg	aac	cca	gaa	act	ttc	aaa	tgg	atg	ctt	aaa	1008
Leu	Pro	Glu	Glu	Met	Arg	Asn	Pro	Glu	Thr	Phe	Lys	Trp	Met	Leu	Lys	
			325					330						335		
aat	cct	cag	tac	cgt	caa	caa	cta	cag	gac	atg	ttg	aat	aat	atg	agt	1056
Asn	Pro	Gln	Tyr	Arg	Gln	Gln	Leu	Gln	Asp	Met	Leu	Asn	Asn	Met	Ser	
		340					345					350				
ggg	agt	ggc	gaa	tgg	gac	aag	cga	atg	aca	gat	aca	ttg	aag	aat	ttt	1104
Gly	Ser	Gly	Glu	Trp	Asp	Lys	Arg	Met	Thr	Asp	Thr	Leu	Lys	Asn	Phe	
	355					360						365				
gac	ctg	aat	agt	cct	gaa	gtg	aag	caa	caa	ttc	aat	caa	ata	gga	cta	1152
Asp	Leu	Asn	Ser	Pro	Glu	Val	Lys	Gln	Gln	Phe	Asn	Gln	Ile	Gly	Leu	
	370					375					380					
act	cca	gaa	gaa	gtc	ata	tct	aag	atc	atg	gag	aac	cct	gat	gtt	gcc	1200
Thr	Pro	Glu	Glu	Val	Ile	Ser	Lys	Ile	Met	Glu	Asn	Pro	Asp	Val	Ala	
385				390					395					400		

atg gca ttc cag aat cct aga gtc caa gca gcg tta atg gaa tgc tca 1248  
Met Ala Phe Gln Asn Pro Arg Val Gln Ala Ala Leu Met Glu Cys Ser  
405 410 415

gag aac cca atg aac atc atg aag tac caa aac gac aaa gag gta atg 1296  
Glu Asn Pro Met Asn Ile Met Lys Tyr Gln Asn Asp Lys Glu Val Met  
420 425 430

gat gtg ttc aac aag ata tcg cag ctc ttc cca gga atg acg ggt tga 1344  
Asp Val Phe Asn Lys Ile Ser Gln Leu Phe Pro Gly Met Thr Gly  
435 440 445

&lt;210&gt; 6

&lt;211&gt; 447

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 6

Met Glu Asn Leu Thr Leu Val Ser Cys Ser Ala Ser Ser Pro Lys Leu  
1 5 10 15  
Leu Ile Gly Cys Asn Phe Thr Ser Ser Leu Lys Asn Pro Thr Gly Phe  
20 25 30  
Ser Arg Arg Thr Pro Asn Ile Val Leu Arg Cys Ser Lys Ile Ser Ala  
35 40 45  
Ser Ala Gln Ser Gln Ser Pro Ser Ser Arg Pro Glu Asn Thr Gly Glu  
50 55 60  
Ile Val Val Val Lys Gln Arg Ser Lys Ala Phe Ala Ser Ile Phe Ser  
65 70 75 80  
Ser Ser Arg Asp Gln Gln Thr Thr Ser Val Ala Ser Pro Ser Val Pro  
85 90 95  
Val Pro Pro Pro Ser Ser Ser Thr Ile Gly Ser Pro Leu Phe Trp Ile  
100 105 110  
Gly Val Gly Val Gly Leu Ser Ala Leu Phe Ser Tyr Val Thr Ser Asn  
115 120 125  
Leu Lys Lys Tyr Ala Met Gln Thr Ala Met Lys Thr Met Met Asn Gln  
130 135 140  
Met Asn Thr Gln Asn Ser Gln Phe Asn Asn Ser Gly Phe Pro Ser Gly  
145 150 155 160  
Ser Pro Phe Pro Phe Pro Phe Pro Pro Gln Thr Ser Pro Ala Ser Ser  
165 170 175  
Pro Phe Gln Ser Gln Ser Gln Ser Ser Gly Ala Thr Val Asp Val Thr  
180 185 190  
Ala Thr Lys Val Glu Thr Pro Pro Ser Thr Lys Pro Lys Pro Thr Pro  
195 200 205  
Ala Lys Asp Ile Glu Val Asp Lys Pro Ser Val Val Leu Glu Ala Ser  
210 215 220  
Lys Glu Lys Lys Glu Glu Lys Asn Tyr Ala Phe Glu Asp Ile Ser Pro  
225 230 235 240  
Glu Glu Thr Thr Lys Glu Ser Pro Phe Ser Asn Tyr Ala Glu Val Ser  
245 250 255  
Glu Thr Asn Ser Pro Lys Glu Thr Arg Leu Phe Glu Asp Val Leu Gln  
260 265 270  
Asn Gly Ala Gly Pro Ala Asn Gly Ala Thr Ala Ser Glu Val Phe Gln  
275 280 285

Ser Leu Gly Gly Gly Lys Gly Gly Pro Gly Leu Ser Val Glu Ala Leu  
 290 295 300  
 Glu Lys Met Met Glu Asp Pro Thr Val Gln Lys Met Val Tyr Pro Tyr  
 305 310 315 320  
 Leu Pro Glu Glu Met Arg Asn Pro Glu Thr Phe Lys Trp Met Leu Lys  
 325 330 335  
 Asn Pro Gln Tyr Arg Gln Gln Leu Gln Asp Met Leu Asn Asn Met Ser  
 340 345 350  
 Gly Ser Gly Glu Trp Asp Lys Arg Met Thr Asp Thr Leu Lys Asn Phe  
 355 360 365  
 Asp Leu Asn Ser Pro Glu Val Lys Gln Gln Phe Asn Gln Ile Gly Leu  
 370 375 380  
 Thr Pro Glu Glu Val Ile Ser Lys Ile Met Glu Asn Pro Asp Val Ala  
 385 390 395 400  
 Met Ala Phe Gln Asn Pro Arg Val Gln Ala Ala Leu Met Glu Cys Ser  
 405 410 415  
 Glu Asn Pro Met Asn Ile Met Lys Tyr Gln Asn Asp Lys Glu Val Met  
 420 425 430  
 Asp Val Phe Asn Lys Ile Ser Gln Leu Phe Pro Gly Met Thr Gly  
 435 440 445

<210> 7  
 <211> 2163  
 <212> DNA  
 <213> Arabidopsis thaliana

<220>  
 <221> CDS  
 <222> (1)..(2163)

<400> 7  
 atg tct agg gag gat ttt agt gat aca ctt cga gta ctt gtt gca act 48  
 Met Ser Arg Glu Asp Phe Ser Asp Thr Leu Arg Val Leu Val Ala Thr  
 1 5 10 15  
 gat tgc cac ttg ggc tac atg gag aag gat gaa att agg cgg cat gat 96  
 Asp Cys His Leu Gly Tyr Met Glu Lys Asp Glu Ile Arg Arg His Asp  
 20 25 30  
 tca ttt aag gct ttc gaa gag ata tgt tct ata gct gag gag aaa cag 144  
 Ser Phe Lys Ala Phe Glu Glu Ile Cys Ser Ile Ala Glu Glu Lys Gln  
 35 40 45  
 gtg gac ttc tta ctc ctc gga ggt gat ctt ttt cat gag aat aaa ccc 192  
 Val Asp Phe Leu Leu Leu Gly Gly Asp Leu Phe His Glu Asn Lys Pro  
 50 55 60  
 tct aga act acg tta gtt aaa gcc att gaa att ctt cgt cgc cac tgt 240  
 Ser Arg Thr Thr Leu Val Lys Ala Ile Glu Ile Leu Arg Arg His Cys  
 65 70 75 80  
 ctg aat gat aaa cca gtg cag ttt caa gta gtc agc gac cag aca gta 288  
 Leu Asn Asp Lys Pro Val Gln Phe Gln Val Val Ser Asp Gln Thr Val

85	90	95	
aat ttt cag aat gcg ttt ggt caa gtc aat tac gag gat cca cac ttc			336
Asn Phe Gln Asn Ala Phe Gly Gln Val Asn Tyr Glu Asp Pro His Phe			
100	105	110	
aat gta ggc ttg ccc gtg ttc agt att cat gga aac cat gat gat cca			384
Asn Val Gly Leu Pro Val Phe Ser Ile His Gly Asn His Asp Asp Pro			
115	120	125	
gcc gga gtg gac aat ctt tct gca att gat att ctt tcc gca tgc aac			432
Ala Gly Val Asp Asn Leu Ser Ala Ile Asp Ile Leu Ser Ala Cys Asn			
130	135	140	
ctt gtg aac tat ttt gga aag atg gtt ctt ggt ggt tct ggt gtt ggc			480
Leu Val Asn Tyr Phe Gly Lys Met Val Leu Gly Gly Ser Gly Val Gly			
145	150	155	160
cag att act ctc tac cct ata ctt atg aag aag ggc tca aca acc gtg			528
Gln Ile Thr Leu Tyr Pro Ile Leu Met Lys Lys Gly Ser Thr Thr Val			
165	170	175	
gct ctc tat ggt tta gga aac atc agg gat gaa cgt ctc aat aga atg			576
Ala Leu Tyr Gly Leu Gly Asn Ile Arg Asp Glu Arg Leu Asn Arg Met			
180	185	190	
ttt cag acc cca cat gct gtc caa tgg atg agg cct gaa gtt caa gaa			624
Phe Gln Thr Pro His Ala Val Gln Trp Met Arg Pro Glu Val Gln Glu			
195	200	205	
gga tgt gat gtt tct gac tgg ttc aac att ctg gtg ctt cat caa aat			672
Gly Cys Asp Val Ser Asp Trp Phe Asn Ile Leu Val Leu His Gln Asn			
210	215	220	
agg gtg aaa tca aac ccc aaa aat gca ata agt gag cac ttt ctt cca			720
Arg Val Lys Ser Asn Pro Lys Asn Ala Ile Ser Glu His Phe Leu Pro			
225	230	235	240
cgt ttc ctc gac ttc att gtg tgg ggc cat gag cat gaa tgc cta atc			768
Arg Phe Leu Asp Phe Ile Val Trp Gly His Glu His Glu Cys Leu Ile			
245	250	255	
gac ccc cag gag gta tct gga atg ggc ttc cac atc aca caa cca gga			816
Asp Pro Gln Glu Val Ser Gly Met Gly Phe His Ile Thr Gln Pro Gly			
260	265	270	
tct tct gtg gca aca tca ctt att gat ggg gaa tcg aag cca aaa cat			864
Ser Ser Val Ala Thr Ser Leu Ile Asp Gly Glu Ser Lys Pro Lys His			
275	280	285	
gtt ctt ctc tta gaa atc aag gga aat caa tat cgt cct acg aag ata			912
Val Leu Leu Leu Glu Ile Lys Gly Asn Gln Tyr Arg Pro Thr Lys Ile			
290	295	300	
cct ttg aca tct gtg agg cct ttt gag tat aca gag att gtt tta aag			960

Pro Leu Thr Ser Val Arg Pro Phe Glu Tyr Thr Glu Ile Val Leu Lys	
305 310 315 320	
gat gaa agt gat att gat ccc aat gat caa aac tca att ctg gaa cac	1008
Asp Glu Ser Asp Ile Asp Pro Asn Asp Gln Asn Ser Ile Leu Glu His	
325 330 335	
ttg gat aaa gtg gtc aga aat cta ata gag aaa gct agc aaa aaa gct	1056
Leu Asp Lys Val Val Arg Asn Leu Ile Glu Lys Ala Ser Lys Lys Ala	
340 345 350	
gtt aac aga tca gag atc aaa ctc cca ttg gtt cga atc aag gta gat	1104
Val Asn Arg Ser Glu Ile Lys Leu Pro Leu Val Arg Ile Lys Val Asp	
355 360 365	
tat tct gga ttt atg acg ata aat cct caa aga ttt gga cag aaa tat	1152
Tyr Ser Gly Phe Met Thr Ile Asn Pro Gln Arg Phe Gly Gln Lys Tyr	
370 375 380	
gtg gga aag gtt gca aat ccc cag gac att ttg ata ttt tcc aag gct	1200
Val Gly Lys Val Ala Asn Pro Gln Asp Ile Leu Ile Phe Ser Lys Ala	
385 390 395 400	
tct aag aag ggt cgg agc gaa gcc aac atc gat gat tct gag cgg ctt	1248
Ser Lys Lys Gly Arg Ser Glu Ala Asn Ile Asp Asp Ser Glu Arg Leu	
405 410 415	
cgt cca gaa gaa ctg aac cag cag aat ata gaa gct tta gta gct gaa	1296
Arg Pro Glu Glu Leu Asn Gln Gln Asn Ile Glu Ala Leu Val Ala Glu	
420 425 430	
agc aac ctg aaa atg gag atc ctt cca gtt aac gat ctg gat gtt gct	1344
Ser Asn Leu Lys Met Glu Ile Leu Pro Val Asn Asp Leu Asp Val Ala	
435 440 445	
ctt cac aat ttt gtg aac aag gat gat aaa cta gcc ttc tac tca tgc	1392
Leu His Asn Phe Val Asn Lys Asp Asp Lys Leu Ala Phe Tyr Ser Cys	
450 455 460	
gtt cag tac aat ctt caa gag act cgt ggt aaa ctt gca aag gat tca	1440
Val Gln Tyr Asn Leu Gln Glu Thr Arg Gly Lys Leu Ala Lys Asp Ser	
465 470 475 480	
gat gcc aag aaa ttt gag gaa gat gac ttg att ctt aaa gtg gga gag	1488
Asp Ala Lys Lys Phe Glu Glu Asp Asp Leu Ile Leu Lys Val Gly Glu	
485 490 495	
tgc tta gag gaa cgc ttg aaa gat agy tcc act cga ccc act ggt tcc	1536
Cys Leu Glu Glu Arg Leu Lys Asp Arg Ser Thr Arg Pro Thr Gly Ser	
500 505 510	
tca cag ttt tta tcc act gga ttg act tca gag aat ttg aca aaa gga	1584
Ser Gln Phe Leu Ser Thr Gly Leu Thr Ser Glu Asn Leu Thr Lys Gly	
515 520 525	

agc agt ggc atc gcg aat gct tcg ttc agt gat gat gaa gac aca act Ser Ser Gly Ile Ala Asn Ala Ser Phe Ser Asp Asp Glu Asp Thr Thr 530 535 540	1632
cag atg tct ggt tta gct cct ccc act aga gga cga aga ggt tca tcc Gln Met Ser Gly Leu Ala Pro Pro Thr Arg Gly Arg Arg Gly Ser Ser 545 550 555 560	1680
act gct aat aca act cgt ggt aga gct aaa gcc cca acc aga gga cga Thr Ala Asn Thr Thr Arg Gly Arg Ala Lys Ala Pro Thr Arg Gly Arg 565 570 575	1728
ggc cgt ggt aag gcc tca agt gcg atg aag caa acc act ctt gat agt Gly Arg Gly Lys Ala Ser Ser Ala Met Lys Gln Thr Thr Leu Asp Ser 580 585 590	1776
tct ctt ggt ttc cgc cag tct caa aga tct gct tcg gct gct gct tca Ser Leu Gly Phe Arg Gln Ser Gln Arg Ser Ala Ser Ala Ala Ala Ser 595 600 605	1824
gct gcc ttc aaa agt gct tcc acc att gga gaa gat gat gta gat tct Ala Ala Phe Lys Ser Ala Ser Thr Ile Gly Glu Asp Asp Val Asp Ser 610 615 620	1872
cct tca agc gaa gaa gtc gag cct gaa gat ttt aac aaa cct gac agc Pro Ser Ser Glu Glu Val Glu Pro Glu Asp Phe Asn Lys Pro Asp Ser 625 630 635 640	1920
agt tcg gag gac gat gag agc act aaa ggc aaa gga cgt aaa aga cca Ser Ser Glu Asp Asp Glu Ser Thr Lys Lys Lys Gly Arg Lys Arg Pro 645 650 655	1968
gct act act aag aga ggc aga ggt aga ggt tct ggg act tca aaa cgt Ala Thr Thr Lys Arg Gly Arg Gly Arg Gly Ser Gly Thr Ser Lys Arg 660 665 670	2016
ggt aga aaa aac gaa agc tct tct tca ctt aat agg cta ctc agt agc Gly Arg Lys Asn Glu Ser Ser Ser Ser Leu Asn Arg Leu Leu Ser Ser 675 680 685	2064
aaa gac gat gac gag gac gaa gat gat gaa gac aga gaa aag aag ctt Lys Asp Asp Asp Glu Asp Glu Asp Asp Glu Asp Arg Glu Lys Lys Leu 690 695 700	2112
aac aaa tct cag cct cgg gtt aca agg aac tat gga gct cta aga aga Asn Lys Ser Gln Pro Arg Val Thr Arg Asn Tyr Gly Ala Leu Arg Arg 705 710 715 720	2160
taa	2163

&lt;210&gt; 8

&lt;211&gt; 720

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 8

```

Met Ser Arg Glu Asp Phe Ser Asp Thr Leu Arg Val Leu Val Ala Thr
 1           5           10           15
Asp Cys His Leu Gly Tyr Met Glu Lys Asp Glu Ile Arg Arg His Asp
           20           25           30
Ser Phe Lys Ala Phe Glu Glu Ile Cys Ser Ile Ala Glu Glu Lys Gln
           35           40           45
Val Asp Phe Leu Leu Leu Gly Gly Asp Leu Phe His Glu Asn Lys Pro
           50           55           60
Ser Arg Thr Thr Leu Val Lys Ala Ile Glu Ile Leu Arg Arg His Cys
           65           70           75           80
Leu Asn Asp Lys Pro Val Gln Phe Gln Val Val Ser Asp Gln Thr Val
           85           90           95
Asn Phe Gln Asn Ala Phe Gly Gln Val Asn Tyr Glu Asp Pro His Phe
           100          105          110
Asn Val Gly Leu Pro Val Phe Ser Ile His Gly Asn His Asp Asp Pro
           115          120          125
Ala Gly Val Asp Asn Leu Ser Ala Ile Asp Ile Leu Ser Ala Cys Asn
           130          135          140
Leu Val Asn Tyr Phe Gly Lys Met Val Leu Gly Gly Ser Gly Val Gly
           145          150          155          160
Gln Ile Thr Leu Tyr Pro Ile Leu Met Lys Lys Gly Ser Thr Thr Val
           165          170          175
Ala Leu Tyr Gly Leu Gly Asn Ile Arg Asp Glu Arg Leu Asn Arg Met
           180          185          190
Phe Gln Thr Pro His Ala Val Gln Trp Met Arg Pro Glu Val Gln Glu
           195          200          205
Gly Cys Asp Val Ser Asp Trp Phe Asn Ile Leu Val Leu His Gln Asn
           210          215          220
Arg Val Lys Ser Asn Pro Lys Asn Ala Ile Ser Glu His Phe Leu Pro
           225          230          235          240
Arg Phe Leu Asp Phe Ile Val Trp Gly His Glu His Glu Cys Leu Ile
           245          250          255
Asp Pro Gln Glu Val Ser Gly Met Gly Phe His Ile Thr Gln Pro Gly
           260          265          270
Ser Ser Val Ala Thr Ser Leu Ile Asp Gly Glu Ser Lys Pro Lys His
           275          280          285
Val Leu Leu Leu Glu Ile Lys Gly Asn Gln Tyr Arg Pro Thr Lys Ile
           290          295          300
Pro Leu Thr Ser Val Arg Pro Phe Glu Tyr Thr Glu Ile Val Leu Lys
           305          310          315          320
Asp Glu Ser Asp Ile Asp Pro Asn Asp Gln Asn Ser Ile Leu Glu His
           325          330          335
Leu Asp Lys Val Val Arg Asn Leu Ile Glu Lys Ala Ser Lys Lys Ala
           340          345          350
Val Asn Arg Ser Glu Ile Lys Leu Pro Leu Val Arg Ile Lys Val Asp
           355          360          365
Tyr Ser Gly Phe Met Thr Ile Asn Pro Gln Arg Phe Gly Gln Lys Tyr
           370          375          380
Val Gly Lys Val Ala Asn Pro Gln Asp Ile Leu Ile Phe Ser Lys Ala
           385          390          395          400
Ser Lys Lys Gly Arg Ser Glu Ala Asn Ile Asp Asp Ser Glu Arg Leu
           405          410          415

```

Arg Pro Glu Glu Leu Asn Gln Gln Asn Ile Glu Ala Leu Val Ala Glu  
                   420                  425                  430  
 Ser Asn Leu Lys Met Glu Ile Leu Pro Val Asn Asp Leu Asp Val Ala  
                   435                  440                  445  
 Leu His Asn Phe Val Asn Lys Asp Asp Lys Leu Ala Phe Tyr Ser Cys  
                   450                  455                  460  
 Val Gln Tyr Asn Leu Gln Glu Thr Arg Gly Lys Leu Ala Lys Asp Ser  
                   465                  470                  475                  480  
 Asp Ala Lys Lys Phe Glu Glu Asp Asp Leu Ile Leu Lys Val Gly Glu  
                   485                  490                  495  
 Cys Leu Glu Glu Arg Leu Lys Asp Arg Ser Thr Arg Pro Thr Gly Ser  
                   500                  505                  510  
 Ser Gln Phe Leu Ser Thr Gly Leu Thr Ser Glu Asn Leu Thr Lys Gly  
                   515                  520                  525  
 Ser Ser Gly Ile Ala Asn Ala Ser Phe Ser Asp Asp Glu Asp Thr Thr  
                   530                  535                  540  
 Gln Met Ser Gly Leu Ala Pro Pro Thr Arg Gly Arg Arg Gly Ser Ser  
                   545                  550                  555                  560  
 Thr Ala Asn Thr Thr Arg Gly Arg Ala Lys Ala Pro Thr Arg Gly Arg  
                   565                  570                  575  
 Gly Arg Gly Lys Ala Ser Ser Ala Met Lys Gln Thr Thr Leu Asp Ser  
                   580                  585                  590  
 Ser Leu Gly Phe Arg Gln Ser Gln Arg Ser Ala Ser Ala Ala Ser  
                   595                  600                  605  
 Ala Ala Phe Lys Ser Ala Ser Thr Ile Gly Glu Asp Asp Val Asp Ser  
                   610                  615                  620  
 Pro Ser Ser Glu Glu Val Glu Pro Glu Asp Phe Asn Lys Pro Asp Ser  
                   625                  630                  635                  640  
 Ser Ser Glu Asp Asp Glu Ser Thr Lys Gly Lys Gly Arg Lys Arg Pro  
                   645                  650                  655  
 Ala Thr Thr Lys Arg Gly Arg Gly Arg Gly Ser Gly Thr Ser Lys Arg  
                   660                  665                  670  
 Gly Arg Lys Asn Glu Ser Ser Ser Ser Leu Asn Arg Leu Leu Ser Ser  
                   675                  680                  685  
 Lys Asp Asp Asp Glu Asp Glu Asp Asp Glu Asp Arg Glu Lys Lys Leu  
                   690                  695                  700  
 Asn Lys Ser Gln Pro Arg Val Thr Arg Asn Tyr Gly Ala Leu Arg Arg  
                   705                  710                  715                  720

&lt;210&gt; 9

&lt;211&gt; 1434

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1434)

&lt;400&gt; 9

atg atg aca tta aac tca cta tct cca gct gaa tcc aaa gct att tct 48  
 Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu Ser Lys Ala Ile Ser  
   1                  5                  10                  15



ttc ttg gat acc tcc agg ttc aat cca atc cct aaa ctc tca ggt ggg	96
Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro Lys Leu Ser Gly Gly	
20 25 30	
ttt agt ttg agg agg agg gat caa ggg aga ggt ttt gga aaa ggt gtt	144
Phe Ser Leu Arg Arg Arg Asp Gln Gly Arg Gly Phe Gly Lys Gly Val	
35 40 45	
aag tgt tca gtg aaa gtg cag cag caa caa caa cct cct cca gca tgg	192
Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln Pro Pro Pro Ala Trp	
50 55 60	
cct ggg aga gct gtt cct gag gcg cct cgt caa tct tgg gat gga cca	240
Pro Gly Arg Ala Val Pro Glu Ala Pro Arg Gln Ser Trp Asp Gly Pro	
65 70 75 80	
aaa ccc atc tct atc gtt gga tct act ggt tcc atc ggc act cag aca	288
Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Ser Ile Gly Thr Gln Thr	
85 90 95	
ttg gat att gtg gct gag aat cct gac aaa ttt aga gtt gtg gct cta	336
Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe Arg Val Val Ala Leu	
100 105 110	
gct gct ggt tcg aat gtt act cta ctt gct gat cag gta agg aga ttt	384
Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp Gln Val Arg Arg Phe	
115 120 125	
aag cct gcg ttg gtt gct gtt aga aac gag tca ctg att aat gag ctt	432
Lys Pro Ala Leu Val Ala Val Arg Asn Glu Ser Leu Ile Asn Glu Leu	
130 135 140	
aaa gag gct tta gct gat ttg gac tat aaa ccc gag att att cca gga	480
Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Pro Glu Ile Ile Pro Gly	
145 150 155 160	
gag cta gga gtg att gag gtt gcc cga cat cct gaa gct gta acc gtt	528
Glu Leu Gly Val Ile Glu Val Ala Arg His Pro Glu Ala Val Thr Val	
165 170 175	
gtt acc gga ata gta ggt tgt gcg gga ctg aag cct acg gtt gct gca	576
Val Thr Gly Ile Val Gly Cys Ala Gly Leu Lys Pro Thr Val Ala Ala	
180 185 190	
att gaa gca gga aag gac att gct ctt gca aac aaa gag aca tta atc	624
Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn Lys Glu Thr Leu Ile	
195 200 205	
gca ggt ggt cct ttc gtg ctt ccg ctt gcc aac aaa cat aat gta aag	672
Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn Lys His Asn Val Lys	
210 215 220	
att ctt ccg gca gat tca gaa cat tct gcc ata ttt cag tgt att caa	720
Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile Phe Gln Cys Ile Gln	
225 230 235 240	

ggt ttg cct gaa ggc gct ctg cgc aag ata atc ttg act gca tct ggt Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile Leu Thr Ala Ser Gly 245 250 255	768
gga gct ttt agg gat tgg cct gtc gaa aag cta aag gaa gtt aaa gta Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu Lys Glu Val Lys Val 260 265 270	816
gcg gat gcg ttg aag cat cca aac tgg aac atg gga aag aaa atc act Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met Gly Lys Lys Ile Thr 275 280 285	864
gtg gac tct gct acg ctt ttc aac aag ggt ctt gag gtc att gaa gcg Val Asp Ser Ala Thr Leu Phe Asn Lys Gly Leu Glu Val Ile Glu Ala 290 295 300	912
cat tat ttg ttt gga gct gag tat gac gat ata gag att gtc att cat His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile Glu Ile Val Ile His 305 310 315 320	960
cct caa agt atc ata cat tcc atg att gaa aca cag gat tca tct gtg Pro Gln Ser Ile Ile His Ser Met Ile Glu Thr Gln Asp Ser Ser Val 325 330 335	1008
ctt gct caa ttg ggt tgg cct gat atg cgt tta ccg att ctc tac acc Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr 340 345 350	1056
atg tca tgg ccc gat aga gtt cct tgt tct gaa gta act tgg cct aga Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Pro Arg 355 360 365	1104
ctt gac ctt tgc aaa ctc ggt tca ttg act ttc aag aaa cca gac aat Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn 370 375 380	1152
gtg aaa tac cca tcc atg gat ctt gct tat gct gct gga cga gct gga Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly 385 390 395 400	1200
ggc aca atg act gga gtt ctc agc gcc gcc aat gag aaa gct gtt gaa Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn Glu Lys Ala Val Glu 405 410 415	1248
atg ttt att gat gaa aag ata agc tat ttg gat atc ttc aag gtt gtg Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp Ile Phe Lys Val Val 420 425 430	1296
gaa tta aca tgc gat aaa cat cga aac gag ttg gta aca tca ccg tct Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu Val Thr Ser Pro Ser 435 440 445	1344
ctt gaa gag att gtt cac tat gac ttg tgg gca cgt gaa tat gcc gcg Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala Arg Glu Tyr Ala Ala	1392

450                      455                      460

gat gtg cag ctt tct tct ggt gct agg cca gtt cat gca tga                      1434

Asp Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala

465                      470                      475

<210> 10  
 <211> 477  
 <212> PRT  
 <213> Arabidopsis thaliana

<400> 10

Met	Met	Thr	Leu	Asn	Ser	Leu	Ser	Pro	Ala	Glu	Ser	Lys	Ala	Ile	Ser
1				5					10					15	
Phe	Leu	Asp	Thr	Ser	Arg	Phe	Asn	Pro	Ile	Pro	Lys	Leu	Ser	Gly	Gly
		20						25					30		
Phe	Ser	Leu	Arg	Arg	Arg	Asp	Gln	Gly	Arg	Gly	Phe	Gly	Lys	Gly	Val
		35					40					45			
Lys	Cys	Ser	Val	Lys	Val	Gln	Gln	Gln	Gln	Pro	Pro	Pro	Ala	Trp	
	50					55				60					
Pro	Gly	Arg	Ala	Val	Pro	Glu	Ala	Pro	Arg	Gln	Ser	Trp	Asp	Gly	Pro
65					70					75				80	
Lys	Pro	Ile	Ser	Ile	Val	Gly	Ser	Thr	Gly	Ser	Ile	Gly	Thr	Gln	Thr
			85						90					95	
Leu	Asp	Ile	Val	Ala	Glu	Asn	Pro	Asp	Lys	Phe	Arg	Val	Val	Ala	Leu
			100					105					110		
Ala	Ala	Gly	Ser	Asn	Val	Thr	Leu	Leu	Ala	Asp	Gln	Val	Arg	Arg	Phe
		115					120					125			
Lys	Pro	Ala	Leu	Val	Ala	Val	Arg	Asn	Glu	Ser	Leu	Ile	Asn	Glu	Leu
	130					135					140				
Lys	Glu	Ala	Leu	Ala	Asp	Leu	Asp	Tyr	Lys	Pro	Glu	Ile	Ile	Pro	Gly
145				150					155					160	
Glu	Leu	Gly	Val	Ile	Glu	Val	Ala	Arg	His	Pro	Glu	Ala	Val	Thr	Val
			165					170						175	
Val	Thr	Gly	Ile	Val	Gly	Cys	Ala	Gly	Leu	Lys	Pro	Thr	Val	Ala	Ala
			180					185					190		
Ile	Glu	Ala	Gly	Lys	Asp	Ile	Ala	Leu	Ala	Asn	Lys	Glu	Thr	Leu	Ile
	195					200						205			
Ala	Gly	Gly	Pro	Phe	Val	Leu	Pro	Leu	Ala	Asn	Lys	His	Asn	Val	Lys
	210					215					220				
Ile	Leu	Pro	Ala	Asp	Ser	Glu	His	Ser	Ala	Ile	Phe	Gln	Cys	Ile	Gln
225					230					235				240	
Gly	Leu	Pro	Glu	Gly	Ala	Leu	Arg	Lys	Ile	Ile	Leu	Thr	Ala	Ser	Gly
			245					250						255	
Gly	Ala	Phe	Arg	Asp	Trp	Pro	Val	Glu	Lys	Leu	Lys	Glu	Val	Lys	Val
		260						265					270		
Ala	Asp	Ala	Leu	Lys	His	Pro	Asn	Trp	Asn	Met	Gly	Lys	Lys	Ile	Thr
		275					280					285			
Val	Asp	Ser	Ala	Thr	Leu	Phe	Asn	Lys	Gly	Leu	Glu	Val	Ile	Glu	Ala
	290					295					300				
His	Tyr	Leu	Phe	Gly	Ala	Glu	Tyr	Asp	Asp	Ile	Glu	Ile	Val	Ile	His
305					310					315				320	
Pro	Gln	Ser	Ile	Ile	His	Ser	Met	Ile	Glu	Thr	Gln	Asp	Ser	Ser	Val
			325						330					335	

Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr  
                   340                  345                  350  
 Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Pro Arg  
                   355                  360                  365  
 Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn  
                   370                  375                  380  
 Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly  
 385                  390                  395                  400  
 Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn Glu Lys Ala Val Glu  
                   405                  410                  415  
 Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp Ile Phe Lys Val Val  
                   420                  425                  430  
 Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu Val Thr Ser Pro Ser  
                   435                  440                  445  
 Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala Arg Glu Tyr Ala Ala  
                   450                  455                  460  
 Asp Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala  
 465                  470                  475

&lt;210&gt; 11

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
           oligonucleotide

&lt;400&gt; 11

gcggacatct acatTTTTga

20

&lt;210&gt; 12

&lt;211&gt; 1353

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 12

gctgggtaag tagatcgttg catcactatg agatgtctaa gcttcttagg gatcaatatg 60  
 acgctgaagg cgcttgatg attatcaaatt ctggatctcc aggcgcaaaa tctcaggtca 120  
 gtttcatcat tctcaaggca cttacagttt ccaactcttt gcttgtaact tagtttctgt 180  
 ttgttcttaa acatattttg aggatttgca gatatggaca gagcaagttg taagtatgta 240  
 tatcaaatgg gcagaaaggc taggcaaaaa cgcgcgggtg gctgagaaat gtagtttatt 300  
 gagtaataaaa agtggcgtaa gttcagccac gatagagttt gaattcgagt ttgcttatgg 360  
 ttatctctta ggtgagcgag gtgtgcaccg ccttatcata agttccactt ctaatgaggt 420  
 atacattata agttataact ctctttctcg taactaatca ctttcgtgtc cattatcatg 480  
 gcccgggaaa gaattaaaag aggttttctt tgcgccagga atgttcagcg actgttgata 540  
 tcataccact attcttgaga gcatctcctg attttgaagt aaaggaaggt gatttgattg 600  
 tatcgtatcc tgcaaaagag gatcacaata tagctgagaa tatgggttgt atccaccata 660  
 ttccgagtg agtaacacta caatcttcag gtattcttga gtgtgttgtt agttgttaca 720  
 ctttggttta ctgcatttta tgcagattat ataactgag gtttttgatg caggagaaag 780  
 aaaccgggtt gcaaacagga tcaaagctct aaaccgggtt aaggcgaagc tacttgtgat 840  
 agcaaaagag caaaagggtt cggatgtaaa taaaatcgac agcaagaaca ttttggaacc 900

```

gcgggaagaa accaggagtt atgtctctaa gggtcacaag atggtggttg atagaaaaac 960
cgggttagag attctggacc tgaaatcggg cttggatgga aacattggac cactccttgg 1020
agctcatatt agcatgagaa gatcaattga tgcgatttag gcttaatcaa ttggtacttt 1080
aattgctttt tgttttgtat ccaaaaagca acaaatgggt gcttgtgtgt gtatatatat 1140
aaccttcttg tccagaacca tatatgattc taaccatcaa acaaagataa gaattggtga 1200
ctatgtgcta tactctacaa tatcaccatg aatacttcaa actagacttt tgataaattt 1260
tgaaacgggt attaccaata aaacgaaaac catgaaactc ttgttttaat tatcagattc 1320
gagaaagttg tgtacaaaca tagctgagaa ggg                                     1353

```

&lt;210&gt; 13

&lt;211&gt; 184

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 13

```

gcttaatcaa ttggtacttt aattgctttt tggtttgtat ccaaaaagca acaaatggkt 60
gcttgtgtgt gtatatatat aaccttcttg gccagaacca tatatgawtc taaccattaa 120
accaagatta gaattggtga ctaaaaaaaaa agaaaaaaaa aaaaaaaaaa aaaaaaaaaa 180
aaaa                                              184

```

&lt;210&gt; 14

&lt;211&gt; 2170

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 14

```

atggttaagcg tttctttaac tctattttct tcattgtttc agttattggc gattgtattc 60
tctgtttatt gtaatcgtat tgtgttaatt ttgatttgac tcactttctc taaagttcaa 120
tttcaaaatt agggattccg agatcataga tattgctttg tttccgagat ttgagttatt 180
cttaagcttg ttttactaac tttcaatatg ttggatttgt tataggcaac tcttgaagat 240
tctttccttg ctgatttgga cgagttatct gacaatgaag cagaattggt gagtgttaaa 300
acacttttga ttactattat ctgtttactt ggaggagcta tgattgtaat tgtagtttgt 360
ttgattatac atatgcagga cgagaatgat ggtgatgttg gaaaggaaga agaagatggt 420
gatatggata tggctgattt agagacactt aactatgatg atctcgataa tgtttctaag 480
ctgcagaaga gtcagagata tgctgatatt atgcataaag tagaggaggc tcttgggaaa 540
gattctgatg gagctgagaa aggaactgtc ttggaagatg atcctgagta taagcttatt 600
gtggattgta atcagctttc ggtcgtatatt gagaatgaaa tcgttattgt ccacaacttt 660
atcaaagaca agtacaagct taagtttcaa gagcttgagt cgttggttca tcaccctatt 720
gactatgcat gtgttggtgaa gaagattggg aatgagacgg atttggctct tgttgatctc 780
gctgaccttc ttccttcagc tattatcatg gttgtttcag ttactgcttt aactacgaaa 840
gggagtgcac tgccagagga tgttttgcaa aagggtgtag aggccttgta tcgggcttta 900
gatccttgatt ccgcaaggaa gaaggtcctt gagtttggtg aaagtaagat gggatctatt 960
gcacctaatc tttctgctat tgttgggagt gctgttgacg ccaaactcat ggggactgct 1020
ggaggtttgt cagcacttgc taaaatgcct gcgtgtaatg ttcaagttct tggccacaag 1080
aggaagaacc ttgctgggtt ttcttctgca acgtctcagt cccgtgtggg ttatctggag 1140
cagacagaga tttaccaaag cagcctcctt ggacttcagg ctgcgctggg caggctcgtg 1200
gctgcaaaat caactttggc agcaagagtt gatgctacta gaggggatcc gttagggata 1260
agtggaaaag ctttcagggg ggagatccgt aagaagattg agaaatggca agaacctcct 1320
cctgcaagac agcctaagcc acttcctgtt cctgattctg aaccgaagaa aagaaggggt 1380
ggtcgcccgtc taagaaaaat gaaagaaagg tagccttttt catcctactt tgtgtcctta 1440
attactgtag attgagttct attcacctgt atttattttg ttgcattctt acgtttctct 1500
ttaaatcagg tatcaagtaa cagatatgag gaagctggcc aacagaatgg cgtttggtac 1560
acctgaagag agctccctcg gtaatatatc ttgtagttac acttgtaaat ggccacttat 1620

```

```

aaggcactta gtctaataatc tactcttcat gatgataggt gatggactag gagaaggtta 1680
tggaatgctt ggccaggcag gaagcaacag gctgcgagta tccagtgttc cgagcaagct 1740
taagattaat gctaaggctg ccaaaaagta agtggtcctc tatttctcct gtgttttttc 1800
ggatttatca tgtaaatatt ttactctta caaattatcc tgccctgttc ttcttccatc 1860
atatctcatt tgogtcttta tatcaattac tttttcaggc tttaaagaaag gcagtatgcy 1920
ggtggtgcga ctacctctgg ttgacatcg agcctggctt tctctcctgt gcaggtacaa 1980
acatttcatt cgattcttga caaaagtgtg atcctgtgtt ccatttgcat cactgtctga 2040
ctccaattgg ttatctatct gacagggaat agagtgtgtc aatcctcagc aggctttagg 2100
attaggaagt gggactcaaa gcacttactt ctcagagtca ggaaccttct cgaagctgaa 2160
gaagatctaa 2170

```

&lt;210&gt; 15

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 15

accttaggcg acttttgaac

20

&lt;210&gt; 16

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 16

aaacgcttac catatctctt tcta

24

&lt;210&gt; 17

&lt;211&gt; 113

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 17

```

aaacactagt cgctcgctgc tcttcaattt tcttctcgaa tctaactgat tgatttctcc 60
ttcgattctt caggagaatc actgaagctt ttgctccca agtagaaaga gat 113

```

&lt;210&gt; 18

&lt;211&gt; 218

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 18

aatatggaag acagagatnc aagtcttgaa aagccgagca ctaaaagtgt aaaaatgaac 60

caaaggtgga aagaaactgc tttctctatc tcatgtctgt ttttaagggtt cttcgggtcac 120  
 ttaagagaca aaaggcattg ttttgatcac tctttggaaa cgttttataa attttatttt 180  
 tgtattagag ccaaaaaaaaa aaaaaaaaaa aaaaaaaaa 218

<210> 19

<211> 4140

<212> DNA

<213> *Arabidopsis thaliana*

<400> 19

cagtacactt agctacactg gatccaagtc tagtgctaaa ctcaaacctc gtgggttttag 60  
 accaaaatct cttcttcttc gtttccttct tcctcatcat atctttcatc ttctccacca 120  
 gaatttggtt taggctctcc ttcttctgtt tctttttctc ccaaagaaac aattagatat 180  
 ggagaacctt accctagttt cttgctcagc ttcttctcca aagctgttaa ttggatgcaa 240  
 tttcacttcc tcgctgaaaa accctactgg gttttctcgt cggactccta atattgtcct 300  
 ccggtgttcc aaaatatctg cctctgctca atctcaatct cctctctcgc gtccggagaa 360  
 cactggagaa atcggttagt ttgcaaattc cactcgacac tctattatag caaatgcca 420  
 aattttccgg aaaaatttcc agtttattac ttttatctat cttattgaaa ctcaaattgc 480  
 gaaccttttt cgactgggtt aatatgagct tatgaattgc tatatctctt aaaaaatcc 540  
 acactttgtg aatttgcaat ttgaattctt gtgaaacca ttcatgttta gaattgttta 600  
 cttaaagttt atgttcgatt tgcagtgggt gtgaaacaga gaagcaaagc ttttgcaagt 660  
 atattttctt cgagtogtga tcaacagaca acttctgttg cttcccttag tgtgcctgtg 720  
 ccaccacctt cttcatcaac catgtaattt tcctgggttt ggacaatgtg cttagtttgt 780  
 atgtcgtttg attcttgggt attaaattgt gttttttctt ttttcttcta gaggatcacc 840  
 acttttcttg attggtgttg gtgttgggtc atcagctttg ttctcatatg tgagtatcaa 900  
 gattccttcc taattttttt ttctctata aatattcttt cttgcttcaa tattgattaa 960  
 taagtgtctg accttttttc ttttctgatg gcattgcagg taacttcaa tttaaaggta 1020  
 cagatacttg gccctctggt ttacaggac ttttgttctc tagtctgttg cagaaccacg 1080  
 attttatgct tcatgtcaac tctagtgtat tgtgctcatg tatctgagat agttttattc 1140  
 actaaactgg ttatcttaac aagggtgaact gtttgtcac acttggtgaa ccgtttatat 1200  
 aagcatcgaa cttttgcctc tctttttttg ggtagtact tgattcgtag atggtaacct 1260  
 acataccatt atgggttttag tgatgcaact cagggtattca gacttatagt cattttcgca 1320  
 actccagtat ttgattgaaa tatattatac aagttgtcat tgctttctct cattattctc 1380  
 taaccggctg ttactctctt tggatttttt tttttgcttt ggttttagaaa tatgcaatgc 1440  
 aaacagctat gaagacgatg atgaacaaa tgaatacgca aaatagccag ttttaataatt 1500  
 ctggattccc atcaggatca ccttttccgt ttccatttcc tcctcaaca agtcctgctt 1560  
 cctcgccatt ccaatctcaa tccagtcctt cagggtgctac cgttgatgtg acagcgacaa 1620  
 aagtagagac acctccttca actaaaccga aacctacacc tgcaaaggat atagaggttg 1680  
 ataagccaag tgttgtctta gaggaagca aagagaagaa agaagaaaag aactatggta 1740  
 gattcttttt ctgtttcaga aatcaacgct ttttcatttg tattctcaat ttgactttc 1800  
 ttctttctc attttccaag cttctaactt ggaagctgat ttacttttgg atgcagcctt 1860  
 tgaagacatt tcaccgagg aaaccacaaa agaaagccca tttagcaact atgcagaagt 1920  
 ctctgaaact aattcccca aagaaactcg cttggttgag gatgtaagtt tcgttttctt 1980  
 ttgtatttcc acagcacacc aagtgggtgat ttaaaaacgt gacatagttt tgctaacctt 2040  
 ctatgctctc ttattgatct ctgggtgaag gtcttgcaaa atggagctgg tccggcaaat 2100  
 ggtgccactg cttcagaggt ttttcaatct ttgggtgagt tattgaattt cagttttcat 2160  
 cactatcagc gcaactgtga tgattcatga ttaaggctac ggatttcaat ttatttttat 2220  
 agcatatgcc aacaattata aacaaaggaa gatatgaaat tgggtataaa gaggaatgag 2280  
 ttggcttcaa aaggatctac tccgttactt ttgtccttct gctagtcgtt gatctgtatt 2340  
 ggtataacca tataagactt gcaggatatt acctgggcaa tctgtttcat atctcatgtg 2400  
 ttatgattct tttttcttat atgctcacgt tattgtctct cttttcctta ttctaaattt 2460  
 aaaactgaat cctgagctct tctattgttt acacaggtgg tgggaaagga gggccgggtt 2520  
 tatctgtaga agctttagag aaaatgatgg aagatccaac agtccagaag atgggtttacc 2580  
 cgtaactcat cttccctagc acattgtctt taaatgcac cattaagttt atctttaaaa 2640

```

ctgggtgctt agtggacatt tggtaacatt gcatgtataa atgcagatac ttgcctgagg 2700
agatgaggaa cccagaaact ttcaaagtga agtcttttaa tattaatcc tgctatcatt 2760
cttttattag tcctcatttt tacatatttc taaagactaa aggttacatg actagctttt 2820
gaatgatgta attcgtttat aggttgatcc aatgggtatc taaatttaaa atacagtttg 2880
gtacttattg tctccgcttg gaattttgta gggatgctta aaaatcctca gtaccgtcaa 2940
caactacagg acatgttgta agagctccat ttacgaaca atttagttgt ttccattgct 3000
tttaagaatg tctaaactat gtaattaaga aatactcttg tttgtttctt ttcatgaatt 3060
taggaataat atgagtggga gtggtgaatg ggacaagcga atgcagata cattgaagaa 3120
ttttgacctg aatagtcctg aagtgaagca acaattcagt aagacaaatc tcagtttgta 3180
ccaagttaat agtacgttaa ataggtctga tactcaatga ttgaatctgt atttgtcaga 3240
tcaaatagga ctaactccag aagaagtcac atctaagatc atggagaacc ctgatgttgc 3300
catggcattc cagaatccta gagtccaagc agcggttaag gaagtacgtt ttcttttaac 3360
ctgaataaga gaattgctta attttacccc acttctttct tcatacaaaa cagaaaccaa 3420
ttacattctt gttgttggtg cagtgtcag agaaccat gaacatcatg aagtaccaa 3480
acgacaaaga ggtaataata ctgccacttc tccattgccc aaaaaggcga ttactttttt 3540
aagaaatttg aggttattat acattgattg caggtaatgg atgtgttcaa caagatctcg 3600
cagctcttcc caggaatgac ggggtgaaaa agctcacgtc tttggttcta tcaaaaatgt 3660
cacattgtct ttagcttttt gtaggagaa aaaaatgtt ttttttttgc aaagagtctt 3720
cagttttggt cagatcagag aattgtgtac catgttaatc ttaaacgcgg tcgggaattg 3780
gagtcgtgtg aaaacgccgc tgctgtgtt tggatgaat attatacaat agaatttgtt 3840
gtcttaccaa aaaaagtcta tgaagacact gaagagcaa ttattatttt taagggaaaa 3900
tttccaaaat aaacttcatt tattcaaaat ttgcttgaaa aaacctcaat ttttttgtt 3960
tgagattgtg tgaataaatc tgccaatatt ttgttttagc aatttaaaaa attgaagt 4020
ttttctcgca aattttaaat agttgtgatt tattttggaa ttttacctta tttttaatat 4080
ccaaaaggag aagtgacgtg gcgatatcga agcgggttaa tgaagtgatg gccccatctt 4140

```

&lt;210&gt; 20

&lt;211&gt; 77

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 20

```

ccacgcgtcc gctccaccag aatttgtttt aggcctctct tcttctgttt cttttcttcc 60
caaagaaaca attagat 77

```

&lt;210&gt; 21

&lt;211&gt; 354

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 21

```

aaaagctcac gtctttggtt ctatcaaaaa tgtcacattg tctttagctt tttgtagggg 60
gaaaaaatg tttttttttt tgcaagagt cttcagtttt ggtcagatca gagaattgtg 120
taccatgtta atcttaaacg cggtcgggaa ttggagtcgt gtgaaaacgc cgctgctgtt 180
gtttggatg aatattatac aatagaattt gttgtcttac caaaaaaagt ctatgaagac 240
actgaagagc aaattattat ttttaaggga aaatttccaa aataaacttc atgtattcaa 300
aatttgcttg aaaaaacctc aatttttttt gttgaaaaaa aaaaaaaaaa aaaa 354

```

&lt;210&gt; 22

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence



&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 22

cagaccacaa taccttcaaa aata

24

&lt;210&gt; 23

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
oligonucleotide

&lt;400&gt; 23

ccattgtgtc tccctcccgc tgtt

24

&lt;210&gt; 24

&lt;211&gt; 5077

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 24

```

atgattgtaa aacttgacag ggaggatttt agtgatacac ttcgagtact tgttgcaact 60
gattgccact tgggctacat ggagaaggat gaaattaggc ggcatgattc atttaaggct 120
ttcgaagaga tatgttctat agctgaggag aaacaggctc ggtattcagt atctatccct 180
tgccagtatt atcttgcggt tgaatcatct aacatattat cttaaataaa aatcttctcc 240
caatattatg agtagtaaac agtgttctac ctaattttaa caaaaattca accaattgcg 300
aggaagaatt ctacagaaagt ttcatatctt cttttttcac tcttttgaaa cagggtggact 360
tcttactcct cggaggtgat ctttttcatg agaataaacc ctctagaact acggttagtta 420
aagccattga aattcttcgt cgccactgtc tgaatgataa accagtgcag tttcaagtag 480
tcagcgacca gacagtaaat ttccagaatg cgtgagactc tatcctttct gctattaatc 540
taatcataac aggaaataat ttcaactgaa ctaattaatt ggcaaattgg ctcaaattcg 600
tgtatagatc tacgtattct tattaatccc ttgacattat tttctggcta cagggttggt 660
caagtcaatt acgaggatcc aacttcaat gtaggcttgc ccgtgttcag tattcatgga 720
aaccatgatg atccagccgg agtggtagat cacttacatc tgcagtctct tgttatgcaa 780
actcatttga ataggtatat agaactggat tagttagtga ataggtattt tattgtgttt 840
ttgttctatg tctcttatgg ctacaggaca atctttctgc aattgatatt ctttccgcat 900
gcaaccttgt gaactatttt ggaaagatgg ttcttggtgg ttctggtgtt ggccagatta 960
ctctctaccc tatacttatg aagaagggtg gtgtaaagaa tttctaacct agacacctgg 1020
ctccccctga cttcttggac tatcatttaa tcaaattaat gtttagggct caacaaccgt 1080
ggctctctat ggtttaggaa acatcaggga tgaacgtctc aatagaatgt ttcaggtaat 1140
ccagaggacc ctacaccttt gctatacaat tgttaattgt gttaatattt attggtttca 1200
cagaccccac atgctgtcca atggatgagg cctgaagttc aagaaggatg tgatgtttct 1260
gactgggtca acattctggt gcttcatcaa aatagggtga ttccattgct ataacatctt 1320
ttagatcggt ttcttactca ttctgtatca gaaatttga tactgtattc atatgacttg 1380
caggggtgaa tcaaacccca aaaatgcaat aagtgaagac tttcttccac gtttcctcga 1440
cttcattgtg tggggccatg agcatgaatg cctaatcgac cccaggtcc atgaaaaatt 1500
tgatttttgg agttattgca tttaaataag agtgagccac aatgttactt gcctctttga 1560
gctaaaagct attaaacttt tgaaggaggt atctggaatg ggcttccaca tcacacaacc 1620

```

```

aggatcttct gtggcaacat cacttattga tggggaatcg aagccaaaac atgttcttct 1680
cttagaaatc aaggttcttc agcaaacaat ctgaaatttc atcttcactt tattcgact 1740
tcattttctg gtcttttttc tctcttttca atcaagcatg taagcttgag tgacttaaaa 1800
tatatgactt acagggaaat caatatcgtc ctacgaagat acctttgaca tctgtgaggc 1860
cttttgagta tacagaggta aagtttactt ttccttaata tgttatgggtg gtggcagact 1920
tctttgctta catattttca aagtgcagat tgttttaaag gatgaaagtg atattgatcc 1980
caatgatcaa aactcaattc tggaaactt ggataaagtg gtacctatc cctcttctca 2040
tagttcatgt ggatatcttt tctcctgccc tttttgaata accagtcact gaatgtctct 2100
actaatatct aaaaaattgt taggtcagaa atctaataga gaaagctagc aaaaaagctg 2160
ttaacagatc agagatcaaa ctcccattgg ttogaatcaa ggtaacttgt ttccaagttt 2220
tcttcaaact gctgcaaatt ctagcaacac tcatataatt aaacctttat tttctaacc 2280
aactctagag gctaggcttt gccagtttga tgcatgcaca cccatagcca caaacagata 2340
attgttatta agaattattaa atgactgaca aaagactaag atctgcttca tctttcaggt 2400
agattattct ggatttatga cgataaatcc tcaaagattt ggacagaaat atgtgggaaa 2460
ggtagctaga aattagttac tgtaacatga tggtcaccat acttctttga atgttggcta 2520
actaatgaca aagtcccaaa cacttacagg ttgcaaatcc ccaggacatt ttgatatttt 2580
ccaaggcttc taagaagggt cggagcgaag gtaagggtcat tgggtgacta gtaatttata 2640
caattttgtt tggattagat tgatgcacgt gcttttactc taacttgtaa tagcttatct 2700
ggcaaaaatt acggttaagt agtgtatctg agatatagta atgtagaaca atatgggcct 2760
atgataacct cctttgttgt tttattgtcg gtattataat tctcgtcata tatatcatga 2820
ctactaactt tctgttgtgt ggagcttgat attgatgtat tgagtgttaa ttttcttct 2880
gttccacttt tcttggtata gttcatgttt ctctgtgtgt aacctatagc atcaaaattt 2940
tgcaaatctt atggattatc tctagttagt atatatgga aatttgccat tttgataatt 3000
tttttgtcta gtgaattgaa tggcaatgat gcatgtcctg atggttgtcc agtgatccag 3060
ttatgatata ttcaatctt ccatttcaca gccaacatcg atgattctga gcggcttcgt 3120
ccagaagaac tgaaccagca gaatatagaa gctttagtag ctgaaagcaa cctggtacat 3180
cctgcaacct tctttcctta tgatttgttt attatcgtca acccctgtag aactttgcca 3240
cagaatgata tagacttggg tagttaccaaa atgggcatga gtacactatg ggatgatcat 3300
tctattttct tccgcagaaa atggagatcc ttccagttaa cgatctggat gttgtctctc 3360
acaattttgt gaacaaggat gataaactag cttctactc atgogttcag tacaatcttc 3420
aagagactcg tgtatgtact attttttact tcaccattca atacaaagt ctgcatagga 3480
tattattttt atttcgtagc acgtccttgt tattgtttt atgatttate tcttccctct 3540
ttttgtacag ggtaaacttg caaaggattc agatgccaa aaatttgagg aagatgactt 3600
gattcttaaa gtgggagagt gcttagaggg aagaagatat agattcagtt agttctgcg 3660
cagattatga gaaccagcag aatattgatc tcaattgcat tattgttcgt gcaggaacgc 3720
ttgaaagata ggtccactcg acccactggg tccacacagt ttttatccac tggattgact 3780
tcagagggtt aaattctctt ttttagattt tccctgcctc tgccttccg ttggtttctc 3840
acagtgtcat tttctacctg agattggtac agaatttgac aaaaggaagc agtggcatcg 3900
cgaatgcttc gttcagtgat gatgaagaca caactcagat gtctggttta gtcctccca 3960
ctagaggacg aagaggttca tccactgcta atacaactcg tggtagagct aaagcccca 4020
ccagaggacg aggcctggtt aaggcctcaa gtgcatgaa gcaaaccact cttgatagtt 4080
ctcttggttt ccgccagtct caaaggtaac tttttgacag cacatttaac cagtttaggg 4140
taggattcac ggacgtgcaa ggaaatgatt ggcataccta gctagctaat gttatgtccc 4200
taatttgtct ttcataagatc tgcttcggct gctgcttcag ctgccttcaa aagtgtctcc 4260
accattggag aagatgatgt agattctcct tcaagcgaag aagtcgagcc tgaagatttt 4320
aacaacactg acagcagttc ggtatggact attccttaca ctgttattca tttgttctc 4380
accataagaa agcccatgta aaaacttgac aacatataac ttttggcatt cttatttctc 4440
tatttgaagt aaattttgog ttttacttt tcttgattct tgtttgatat ccactaaagg 4500
aggacgatga gagcactaaa ggcaaaggac gtaaaagacc agctactact aagagaggca 4560
gaggtagagg ttctgggact tcaaacgtg gtagaaaaaa cgaaagctct tcttactta 4620
ataggctact cagtagcaaa gacgatgacg aggcgaaga tgatgaagac agagaaaaga 4680
agcttaacaa atctcagcct cgggtttgtt aatcacatct attttccctt ctttcgctgc 4740
ttattagcag gtttttagtaa gttgttggtt accatttgag atcaaagctc acttaatagt 4800
acaatttgaa tatgcagggt acaaggaact atggagctct aagaagataa atacatatca 4860
aaccctaact tctgacatca caacgaagct tcatttttct gttattttct agcgacctct 4920

```

caagcgggaac aacttctgaa gaagagaaat tagtactaac aagagttctg tgagatgatg 4980  
 tacagagaat tttgtagtgt ttttttttct tgctcttttt aagggttacgt tgttgatgaa 5040  
 tgaggcaata tgattaacgt cagtaagaag tctaaaa 5077

<210> 25

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
 oligonucleotide

<400> 25

tgtaaaacga cggccagt

18

<210> 26

<211> 255

<212> DNA

<213> Arabidopsis thaliana

<400> 26

atacatatca aaccccaatc tctgacatca caacgaagct tcatttttct gttattttct 60  
 agcgacctct caagcgggaac aacttctgaa gaagagaaat tagtactaac aagagttctg 120  
 tgagatgatg tacagagaat tttgtagtgt ttttttttct tgctcttttt aagggttacgt 180  
 tgttgatgaa tgaggcaata tgattaacgt cagtaagaag tctaaaaaaaa aaaaaaaaaa 240  
 aaaaaaaaaa aaaaaa 255

<210> 27

<211> 2935

<212> DNA

<213> Arabidopsis thaliana

<400> 27

tcatgcatga actggcctag caccagaaga aagctgcaca ttcgcggcat attcacgtgc 60  
 ccacaagtca tagtgaacaa tctcttcaag agacgggtgat gttaccaact cgtttcgtatg 120  
 tttatcgcat gttaattcca caaccttgaa gatatccaaa tagcttatcc tgtaaacaaa 180  
 agtgagaata taacaattg tgattcgtat caagaacttc attgagatgc tcaaaactga 240  
 aaaataattc ttacttttca tcaatgaaca tttcaacagc tttctcattg gcggcgctga 300  
 gaactccagt cattgtgcct ccagctcgtc cagcagcata agcaagatcc atggatgggt 360  
 atttcacatt gtctgggttc ttgaaagtca atgaaccgag tctgccaaaa tccacaattg 420  
 taaacaactt ttggtttttag gtgctgaatg ctgatagata aggcagtgggt cctaaccag 480  
 ttttaactgat ccacaccaa acagtagcaa aataaccaat tgcaaaacca aaccgaagac 540  
 cgattcgggt tcatttttta tcttatctaa acaacctaaa accaaactga aaacaagatt 600  
 ggggaacttt tcttgggtgat aattaaaatt ttcaactaag cttagcttca cacttgataa 660  
 acagagagta tataaatgtg gttagcttac ttgcaaaggt caagtcttgg ccaagttact 720  
 tcagaacaag gaactctatc gggccatgac atgggtgtaga gaatcggtaa acgcatatca 780  
 ggccaaccca attgagcaag cacagatgaa tcctgtggaa caaaacaaat acatgttata 840  
 cagttatttt tttaaaacgg gaaaaataat aatttagtta gtaatgtttc agcaagacct 900  
 gtgtttcaat catggaatgt atgatacttt gcggatgaat gacaatctct atatcgctcat 960  
 actcagctcc aaacaaataa tgcgcttcaa tgacctcaag accctgtttc aaaaaatcaa 1020  
 gaactcatct accttgatca aaggtatttt caaaatcaga gtttaacctt aggagaaaat 1080

```

aatcttaacc ttgttgaaaa gcgtagcaga gtccacagtg attttctttc ccatgttcca 1140
gtttggatgc ttcaacgcat ccgctacttt aacttctctt agctttttcga caggccaatc 1200
cctttttcaa aatccagtga aaagtttcca ttaaccaaac gagaattgag aagaaaaaaa 1260
gtctatgcag agagagaaga atatcgaaac aaacctaaaa gctccaccag atgcagtcaa 1320
gattatcttg cgcagagcgc cttcaggcaa accttgaata cactagagaa cataaaaaga 1380
gattttttcac tcaaattgcc agaggttgaa cttgcattaa gaccaacgct gaactcaata 1440
tgaaagttga ggtacttaat tctatgtgat ttgtgatacc tgaaatatgg cagaatgttc 1500
tgaatctgcc ggaagaatct ttacattatg tttgttgcca agcgggaagca cgaaaggacc 1560
acctgcgatt aatgtctctt tgtttgcaag agcaatgtcc tttcctgctt caattgcagc 1620
aaccgtaggc tgcagtaaaa ataagcaaca agctttatca tctgcaactt tcttttttca 1680
tatcctctta ataaggttta ataacaaaaa attagagtat atacctttag tcccgcacaa 1740
cctactattc cggtacaacac ggttacagct tcaggatgtc gggcaacctg ttgatgaaca 1800
taataagtaa aaacctatct aactacaat caaaactaac aaatgaacta acctcaatca 1860
ctccttgctc tcttggataa atctcgagtt tatagtccaa atcagctaaa gcctctttaa 1920
gctcattaat cagtgaactcg tttctaacag caaccaatgc aggcctaaat ctccttacct 1980
gccaccattc aaaatagaat cacagaacca tactatagag atttcttgag attgcagaag 2040
caaaagccta aaccagaacc tgatttctct ggtttgatct gatacataac gaggtaatac 2100
tatcttgctt atgatactac cactgaactg agaattaaac tgaattccaa gtggtctgaa 2160
tgacaaattg gagagactca atactaattt ttttacaat gaagccaact tacctgatca 2220
gcaagtagag taacattcga accagcagct agagccacaa ctctgaattt gtcaggattc 2280
tcagccacaa tatccaatgt ctgcaaaatg gaagttcttg tcgataaaaa tgatgcaaca 2340
ataactcagt aagaaaaaaa tatcattctt ctatgagctt agtcattcat aagacaaact 2400
taaagtctgg tcatactcaa gaactgcaca ataatgcctt aatcgaaata aaacctgagt 2460
gccaatagaa ccagtagatc caacgataga gatgggtttt ggtccatccc aagattgacg 2520
aggcgcctca gggacagctc tcccaggcca tgctggagga ggttgttgtt gctgctgcac 2580
tttactgaa cacttaacac cttttccaaa acctctccct tgattcctcc tcctcaaaact 2640
aaacccacct gtgaaacact ccaaagatgt aaaatttaaa actctacgac cttaaagcaaa 2700
ccaaaaaaa tcgaattgaa gaaataacag attacctaga tagagaaatt cacaagagcc 2760
taagacaact aatgaaagtt tgcaacttta atcgaaaaga gatttgacca aggaggagga 2820
aagaagagag gaagaagaag aaacctgaga gtttagggat tggattgaac ctggagggtat 2880
ccaagaaaga aatagctttg gattcagctg gagatagtga gtttaatgtc atcat 2935

```

&lt;210&gt; 28

&lt;211&gt; 1434

&lt;212&gt; DNA

&lt;213&gt; Arabidopsis thaliana

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1434)

&lt;400&gt; 28

```

atg atg aca tta aac tca cta tct cca gct gaa tcc aaa gct att tct 48
Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu Ser Lys Ala Ile Ser
1 5 10 15

```

```

ttc ttg gat acc tcc agg ttc aat cca atc cct aaa ctc tca ggt ggg 96
Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro Lys Leu Ser Gly Gly
20 25 30

```

```

ttt agt ttg agg agg agg rat caa ggg aga ggt ttt gga aaa ggt gtt 144
Phe Ser Leu Arg Arg Arg Xaa Gln Gly Arg Gly Phe Gly Lys Gly Val
35 40 45

```

aag tgt tca gtg aaa gtg cag cag caa caa caa cct cct cca gca tgg	192
Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln Pro Pro Pro Ala Trp	
50 55 60	
cct ggg aga gct gty cct gag gcg cct cgt caa tct tgg gat gga cca	240
Pro Gly Arg Ala Xaa Pro Glu Ala Pro Arg Gln Ser Trp Asp Gly Pro	
65 70 75 80	
aaa ccc atc tct atc gtt gga tct act ggt tcy aty ggc act cag aca	288
Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Xaa Xaa Gly Thr Gln Thr	
85 90 95	
ttg gat att gtg gct gag aat cct gac aaa tty aga gtt gtg gct cta	336
Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe Arg Val Val Ala Leu	
100 105 110	
gct gct ggt tcg aat gtt act cta ctt gct gat cag gta agg aga ttt	384
Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp Gln Val Arg Arg Phe	
115 120 125	
aag cct gcr ttg gtt gct gtt aga aac gag tca ctg att aat gag ctt	432
Lys Pro Xaa Leu Val Ala Val Arg Asn Glu Ser Leu Ile Asn Glu Leu	
130 135 140	
aaa gag gct tta gct gat ttg gac tat aaa cyc gag att att cca gga	480
Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Xaa Glu Ile Ile Pro Gly	
145 150 155 160	
gag cwa gga gtg att gag gtt gcc cga cat cct gaa gct gta acc gtt	528
Glu Xaa Gly Val Ile Glu Val Ala Arg His Pro Glu Ala Val Thr Val	
165 170 175	
gtt acc gga ata gta ggt tgt gcg gga ctg mag cct acg gtt gct gca	576
Val Thr Gly Ile Val Gly Cys Ala Gly Leu Xaa Pro Thr Val Ala Ala	
180 185 190	
att gaa gca gga aag gac att gct ctt gca aac aaa gag aca tta atc	624
Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn Lys Glu Thr Leu Ile	
195 200 205	
gca ggt ggt cct ttc gtg ctt ccg ctt gcc aac aaa cat aat gta aag	672
Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn Lys His Asn Val Lys	
210 215 220	
att ctt ccg gca gat tca gaa cat tct gcc ata ttt cag tgt att caa	720
Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile Phe Gln Cys Ile Gln	
225 230 235 240	
ggt ttg cct gaa ggc gct ctg cgc aag ata atc ttg act gca tct ggt	768
Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile Leu Thr Ala Ser Gly	
245 250 255	
gga gct ttt agg gat tgg cct gtc gaa aag cta aag gaa gtt aaa gta	816
Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu Lys Glu Val Lys Val	
260 265 270	

gcg gat gcg ttg aag cat cca aac tgg aac atg gga aag aaa atc act Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met Gly Lys Lys Ile Thr 275 280 285	864
gtg gac tct gct acg ctt ttc aac aag ggt ctt gag gtc att gaa gcg Val Asp Ser Ala Thr Leu Phe Asn Lys Gly Leu Glu Val Ile Glu Ala 290 295 300	912
cat tat ttg ttt gga gct gag tat gac gat ata gag att gtc att cat His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile Glu Ile Val Ile His 305 310 315 320	960
ckk caa agt atc ata cat tcc atg att gaa aca cag gat tca tct gtg Xaa Gln Ser Ile Ile His Ser Met Ile Glu Thr Gln Asp Ser Ser Val 325 330 335	1008
ctt gct caa ttg ggt tgg cct gat atg cgt tta ccg att ctc tac acc Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr 340 345 350	1056
atg tca tgg ccc gat aga gtt cct tgt tct gaa gta act tgg ccw aga Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Xaa Arg 355 360 365	1104
ctt gac ctt tgc aaa ctc ggt tca ttg act ttc aag aaa cca gac aat Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn 370 375 380	1152
gtg aaa tac cca tcc atg gat ctt gct tat gct gct gga cga gct gga Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly 385 390 395 400	1200
ggc aca atg act gga gtt ctc agc gcc gcc aat gag aaa gct gtt gaa Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn Glu Lys Ala Val Glu 405 410 415	1248
atg tty att gat gaa aag ata agc tat ttg gat atc ttc aag gtt gtg Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp Ile Phe Lys Val Val 420 425 430	1296
gaa tta aca tgc gat aaa cat cga aac gag ttg gta aca tca ccg tct Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu Val Thr Ser Pro Ser 435 440 445	1344
ctt gaa gag att gtt cac tat gac ttg tgg gca cgt gaa tat gcc gcg Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala Arg Glu Tyr Ala Ala 450 455 460	1392
rat gtg cag ctt tct tct ggt gct agg cca gtt cat gca tga Xaa Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala 465 470 475	1434

&lt;211&gt; 477

&lt;212&gt; PRT

&lt;213&gt; Arabidopsis thaliana

&lt;400&gt; 29

```

Met Met Thr Leu Asn Ser Leu Ser Pro Ala Glu Ser Lys Ala Ile Ser
 1          5          10          15
Phe Leu Asp Thr Ser Arg Phe Asn Pro Ile Pro Lys Leu Ser Gly Gly
 20          25          30
Phe Ser Leu Arg Arg Arg Xaa Gln Gly Arg Gly Phe Gly Lys Gly Val
 35          40          45
Lys Cys Ser Val Lys Val Gln Gln Gln Gln Gln Pro Pro Pro Ala Trp
 50          55          60
Pro Gly Arg Ala Xaa Pro Glu Ala Pro Arg Gln Ser Trp Asp Gly Pro
 65          70          75          80
Lys Pro Ile Ser Ile Val Gly Ser Thr Gly Xaa Xaa Gly Thr Gln Thr
 85          90          95
Leu Asp Ile Val Ala Glu Asn Pro Asp Lys Phe Arg Val Val Ala Leu
100          105          110
Ala Ala Gly Ser Asn Val Thr Leu Leu Ala Asp Gln Val Arg Arg Phe
115          120          125
Lys Pro Xaa Leu Val Ala Val Arg Asn Glu Ser Leu Ile Asn Glu Leu
130          135          140
Lys Glu Ala Leu Ala Asp Leu Asp Tyr Lys Xaa Glu Ile Ile Pro Gly
145          150          155          160
Glu Xaa Gly Val Ile Glu Val Ala Arg His Pro Glu Ala Val Thr Val
165          170          175
Val Thr Gly Ile Val Gly Cys Ala Gly Leu Xaa Pro Thr Val Ala Ala
180          185          190
Ile Glu Ala Gly Lys Asp Ile Ala Leu Ala Asn Lys Glu Thr Leu Ile
195          200          205
Ala Gly Gly Pro Phe Val Leu Pro Leu Ala Asn Lys His Asn Val Lys
210          215          220
Ile Leu Pro Ala Asp Ser Glu His Ser Ala Ile Phe Gln Cys Ile Gln
225          230          235          240
Gly Leu Pro Glu Gly Ala Leu Arg Lys Ile Ile Leu Thr Ala Ser Gly
245          250          255
Gly Ala Phe Arg Asp Trp Pro Val Glu Lys Leu Lys Glu Val Lys Val
260          265          270
Ala Asp Ala Leu Lys His Pro Asn Trp Asn Met Gly Lys Lys Ile Thr
275          280          285
Val Asp Ser Ala Thr Leu Phe Asn Lys Gly Leu Glu Val Ile Glu Ala
290          295          300
His Tyr Leu Phe Gly Ala Glu Tyr Asp Asp Ile Glu Ile Val Ile His
305          310          315          320
Xaa Gln Ser Ile Ile His Ser Met Ile Glu Thr Gln Asp Ser Ser Val
325          330          335
Leu Ala Gln Leu Gly Trp Pro Asp Met Arg Leu Pro Ile Leu Tyr Thr
340          345          350
Met Ser Trp Pro Asp Arg Val Pro Cys Ser Glu Val Thr Trp Xaa Arg
355          360          365
Leu Asp Leu Cys Lys Leu Gly Ser Leu Thr Phe Lys Lys Pro Asp Asn
370          375          380
Val Lys Tyr Pro Ser Met Asp Leu Ala Tyr Ala Ala Gly Arg Ala Gly
385          390          395          400

```

Gly Thr Met Thr Gly Val Leu Ser Ala Ala Asn Glu Lys Ala Val Glu  
                    405                    410                    415  
Met Phe Ile Asp Glu Lys Ile Ser Tyr Leu Asp Ile Phe Lys Val Val  
                    420                    425                    430  
Glu Leu Thr Cys Asp Lys His Arg Asn Glu Leu Val Thr Ser Pro Ser  
                    435                    440                    445  
Leu Glu Glu Ile Val His Tyr Asp Leu Trp Ala Arg Glu Tyr Ala Ala  
                    450                    455                    460  
Xaa Val Gln Leu Ser Ser Gly Ala Arg Pro Val His Ala  
465                    470                    475